	- THE CANADIAN ENTOMOLOGIST.	115
	Mesonotum not strongly punctured, and with distinct stripes	
12.	Head produced anteriorly; third joint of antenna at leas times as long at base. *Parectecephala Be Head not distinctly produced; third joint of antenna but longer than broad; dull yellow and black, robust sp with black palpi. *Anthracophaga I	t 1¼ ecker. little
	Hind tibia with a distinct, curved, apical thorn on the and side	erior
14.	Frons with distinct orbital setulæ	17
	Orbits with, at most, weak hairs.	15
15.	Scutellum elongated, disk flattened; male proboscis with long apical hairs	two
	normalPseudohippelates Mal	loch.
	Arista flattened, strap-like Ceratobarys Coqui	llett.
17. 5	Second vein exceptionally short, third costal division in longer than second	ani.
	Arista either broadened and strap-like, or with distinct pu	bes-
P	Arista neither thickened nor distinctly pubescent	00
a a	Arista more or less distinctly flattened, the pubescence of close, giving the arista a knife-like appearance even when rista itself is but slightly flattened	the
.0. 3	cutellum subtriangular, the marginal bristles situated small warts	on
	Warts Melanacheta D	lear
21. S	cutellum elongated, the disk flattened	new

<sup>\*</sup>The line of demarcation between these genera is very unsatisfactory, and though I am rather inclined to reject *Parectecephala* as entitled to generic rank, I have not sufficient material before me to enable me to decide the matter satisfactorily.