

abscess in the left arm. The case eventually made a good recovery. I am inclined to think it had no relation to the operation for the removal of tonsils and adenoids.

Probably no operation in surgery is so abused by the unskilful and also unnecessary removal of tonsils and adenoids. But when properly done and in suitable cases, the results are most satisfactory. The child's intellect, hearing and bodily welfare are preserved.

List of Instruments Used.

7½" dissecting forceps.

Doyen's gag.

St. Clair Thomson's adenoid curette.

Heath's tonsillotome.

Farlow's snare.

Mixter's artery forceps.

Leland's tonsil knives.

Lowenberg's post nasal forceps.

Author's tongue depressing forceps.

Tilley's raspatory for tonsil resector.

Andrew's volsellum.

142 Carlton St., Toronto.

THE ACTION OF ARSENO-PHENYL-GLYCIN UPON *TRYPANOSOMA BRUCEI*.

BY

R. P. CAMPBELL, B.A., M.D., and JOHN L. TODD, M.D.

From the Laboratories, Macdonald College, Ste. Anne de Bellevue, Quebec.

I.

INTRODUCTION.

For two excellent reasons the practitioner in temperate climates must be interested in the experimental work which has been done on the treatment of the various diseases produced by trypanosomes. (1) The first of these reasons is that these researches have given him a third drug, to supplement mercury and potassium iodide in the treatment of syphilis. The second reason is, that during these researches some light has been thrown upon the method of action on the parasites of some of the trypanocides employed. As Ehrlich says, (2) through the continuation of studies such as these on the interdependence of chemical constitution and physiological, or therapeutic, action rational methods of treatment