

blood. In three cases—Experiments 80, 92, and 103—rhythmical movements of the diaphragm were noticed after the heart had ceased beating and after the chest had been opened. It is remarkable that in two of these cases the splanchnic nerve had been divided. The third was a case in which chloroform had been injected into the jugular vein, and in this case there was a synchronous movement of the jaw as well. In all, death and stoppage of the heart had occurred gradually, and in Experiment 103 the heart was still irritable. These movements cannot be called respiration; though the last gasp of a dying animal, that ineffective jerk of the diaphragm, which is such a fatal symptom, is very likely in many cases a movement of the same character. Similar movements, which were continued much longer, occurred in Experiment 104, after the thorax was opened, while the heart was still beating. Still more remarkable convulsions of the muscles of jaws, ears, and fore-feet occurred in Experiment 167, in the case of a dog that had been poisoned with nicotine. These movements continued at regular intervals for more than ten minutes after death, and were sufficiently forceable to jerk the handles of a pressure forceps fixed on the end of the tongue off the table at each spasm. In a rabbit, in Experiment 153, the auricles of the heart continued to beat rhythmically for three hours after it was supposed to be dead from chloroform and its thorax had been laid open. Irritability of the heart after death was noticed in many cases, but seemed to be most marked in cases where ether had been used.

(30) Chloroform injected into the heart through the jugular vein did not cause clotting of the blood, as was the case when ether was injected.

(31) In the course of the experiments of the Committee various drugs were administered in order to ascertain if they had any effect in modifying the action of chloroform. The result showed that none of them had any effect in preventing the typical descent of the blood pressure that occurs when chloroform is inhaled. Atropine, when given in a dose sufficient to paralyse the vagi, of course prevents the action of those nerves in asphyxia, and by increasing the action of the heart it appears to cause a more rapid descent in the blood pressure when chloroform is inhaled, as has been already explained. Morphine appeared in Experiment 162 to render the rise in blood pressure that occurred when the chloroform was discontinued slower and less complete, and to bring about a more or less permanent condition of anaesthesia. It may be noted that the animal used in this experiment was a monkey; and in other experiments with monkeys, when no morphine had been given, it was remarked that the animal, after a few inhalations of chloroform, would often lie quite quiet in a state of semi-in-

sensibility for a long time without further inhalations; still this condition was much more marked in Experiment 162 than in any of the others. No action of this kind was noticed in the dog in Experiment 178, but other experiments (90 and 94) showed that pariah dogs are very indifferent to action of morphine, and it is probable that the dose of morphine in this case was insufficient to bring about the condition noted in the monkey. The peculiar behavior of the heart in Experiment 178 was not the result of the previous administration of morphine, for a similar phenomenon had occurred in other cases (49 and 60) in which no morphine had been given. Experiments 162 and 178 prove conclusively that morphine has no effect in shortening the period that may be allowed to elapse between the cessation of natural respiration and the commencement of artificial respiration.

(32) The other drugs used had no effect upon the action of chloroform except when their own special action became the leading feature in the case—as, for instance, during the vomiting from apomorphine (Experiment 104, Fick 9) or the convulsions produced by nicotine (Experiment 167).

(33) In order to test the alleged danger from shock during chloroform administration, the Committee performed a very large number of those operations which are reputed to be particularly dangerous in this connection—such as extraction of teeth, evulsion of nails, section of the muscles of the eye, snipping of the skin of the anus, etc. In many cases the operation was performed when the animal was merely stupefied by the chloroform and not fully insensible. In such cases a slight variation in the blood pressure would sometimes occur, such as one would expect from the irritation of a sensory nerve or from the struggling that ensued, but in no case in any stage of anaesthesia was there anything even suggestive of syncope or failure of the heart's action. In thrusting a needle into the heart, there was often a momentary but well-marked fall of blood pressure; but even this was absent in all other injuries. If chloroform really had any power to increase the tendency to shock in operations, it is impossible to believe that it would not have been manifest, to some degree at least, in one or other of these numerous experiments. The Commission was, however, not content with this negative result, and determined to ascertain the effect of direct irritation of the vagi during continued chloroform administration. The result of such experiments (65, 117, and others) proved that inhibition of the heart's action prevented, rather than assisted, the fatal effects of prolonged chloroform inhalation. An animal that was put into a condition of extreme danger (from which it could only be restored by means of artificial respiration) by inhalation of chloroform for one minute recovered spontaneously and readily after five minutes of