

LEDA JACKSONI, Gould.

Henley Harbor and Temple Bay, in 10 to 15 fathoms.

LEDA MINUTA, (Mull.) Møller.

Found with the preceding.

MYTILUS EDULIS, Linné.


Henley Harbor, Temple Bay, Fox Harbor, Dead Island, in 1 to 10 fathoms.

MODIOLARIA NIGRA, (Gray) Loven.

Several young specimens were found at L'anse au Loup, 10 fathoms.

MODIOLARIA DISCORS, (Linné) Loven.

Dead Island, near Square Island, in 1 to 4 fathoms.



MODIOLARIA CORRUGATA, (Stimpson) Mørch.

Henley Harbor, in 3 to 7 fathoms.

CRENELLA FABA, (Mull.) (= *C. pectinula* Gld. ?).

Mytilus faba Fabricius, Fauna Grælandica, p. 419.

Fig. 3.—*Crenella faba*, x 4.
From Labrador.

The young of this species were found at L'anse au Loup, 10 to 15 fathoms, sand, mud, kelp; Fox Harbor, Saint Lewis Sound, 1 to 4 fathoms, sand, abundant; Dead Island, near Square Island, shallow water.

CRENELLA GLANDULA, (Totten) H. & A. Ad.

L'anse au Loup, in 8 to 10 fathoms.

CRENELLA DECUSSATA, (Mont.) Macg.

Crenella decussata Verrill, Cat. Mar. Mollusca, p. 578, pl. 41, fig. 7.

Several specimens were found at L'anse au Loup, 10 fathoms; Fox Harbor, Saint Lewis Sound, 3 fathoms.

[THE CUT OF THIS SPECIES WAS MISPLACED AND NOT FOUND AT TIME OF GOING TO PRESS]

PECTEN ISLANDICUS, Møller.

Dead Island, near Square Island, in 1 to 4 fathoms.

ANOMIA ACULEATA, Mull.

L'anse au Loup, 8 to 15 fathoms; Henley Harbor, 10 to 15 fathoms.

RHYN
He
Dead

List

Be
Fu
Fu
Be
? N
L



Fig.
lis.
Lab.
A. S
Fig
rior
S