

*ENOTHERA MICRANTHA*, Horn.

On ballast heaps at Nanaimo, Vancouver Island, 1893. (*John Macoun*, Herb. No. 249.) Introduced from the south. New to Canada.

*PHACELIA HISPIDA*, Gray.

On ballast heaps, Nanaimo, Vancouver Island, 1893. (*John Macoun*, Herb. No. 654.) Introduced from the south. New to Canada.

*KRYNITZIA AMBIGUA*, Gray.

On ballast heaps at Nanaimo, Vancouver Island, 1893. (*John Macoun*, Herb. No. 672.) Introduced from California. New to Canada.

*AM SINCKIA BARBATA*, Greene, *Erythraea*, Vol. II., No. 12, p. 192.

Stout and coarse, erect or decumbent, the branches loosely floriferous throughout, all excepting the uppermost pedicels subtended each by a broad ovate-lanceolate amplexicaul foliaceous bract; sepals 4 or 5 lines long, nearly linear, without rufous or fulvous pubescence, but densely white-hirsute along the margins, sparsely hispid with whitish bristles on the back; corolla small; nutlets ovate-acuminate, closely muricate-tuberculate, without transverse rugosities, but with an elevated and toothed dorsal ridge.

Collected at Cameron Lake, Vancouver Island, 15th July, 1887, by John Macoun. Type specimen in the herbarium of the British Museum. Distributed by Prof. Macoun as *Amsinckia lycopoides*.

*ZANNICHELIA PALUSTRIS*, Linn.

In the Spullamacheen River at Enderby, B.C., 1889. (*Jas. M. Macoun*.) Not before recorded west of the Saskatchewan.