

A rare summer visitor, occasionally straggling here from the prairie sloughs, where they are abundant.

19. Bonaparte's Gull.

(*Larus philadelphia.*)

A common spring and autumn visitor to the great lakes, arriving in April and departing at the end of November. It is probable that some breed in the northern part of the province.

SUBFAMILY STERNINÆ. TERNS.

GENUS STERNA.

SUBGENUS THALASSEUS.

20. Caspian Tern.

(*Sterna caspia.*)

A regular but not common spring and autumn visitor to the great lakes.

SUBGENUS ACTOCHELIDON.

***21. Cabot's Tern.**

(*Sterna sandvicensis aculeiflvida.*)

The capture of a specimen of this tern in the spring of 1882, at Lucknow, is recorded by Mr. McIlwraith.

SUBGENUS STERNA.

22. Forster's Tern.

(*Sterna forsteri.*)

An uncommon summer visitor, breeds throughout its range.

23. Common Tern

(*Sterna hirundo.*)

A regular summer visitor, more frequently seen in spring and autumn.

SUBGENUS STERNULA.

24. Least Tern.

(*Sterna antillarum.*)

An accidental visitor.

GENUS HYDROCHELIDON.

25. Black Tern.

(*Hydrochelidon nigra surinamensis.*)

This Tern is common in some of the large marshes throughout the Province during the summer, but it is not generally distributed. Breeds where it occurs.