THE CANADIAN ENTOMOLOGIST.

marked by a spot on costa, with an outward subcostal angle and an inward one on vein I. Orbicular round, small, black ; reniform a large, diffuse, black cloud. Median shade close to t.-p. line, broad, diffuse, obsolete above. T.-p. line crenulate, inflexed at costa to a spot above reniform and bent inward below cell evenly. A light space beyond it is succeeded by a broad smoky marginal shade, which a faint, pale, subterminal line bisects. The shade is gathered to an ill-defined dark patch on anal angle. Hind wings whitish at base, brownish outwardly, subpellucid. Expanse 20 mm. The wings are not very broad.

Described from one specimen, d, Revelstoke, B. C.

U. S. National Museum, type No. 7325.

Caradrina nitens, n. sp.

Large, slender, close to *mirauda*, Grote, but much blacker. Head and thorax black, not lighter in front; fore wings shining brown-black, orbicular a black dot, reniform a white lunate spot preceded by a black dot; other marks lost, the lines barely traceable in the best specimen. Hind wings silky whitish, smoky along costa and outer edge. Expanse 30 mm.

Described from 8 specimens, Kaslo, B. C., and Turtle Mts., North Dakota (A. H. Verrill).

U. S. National Museum, type No. 7336.

Hadena maida, n. sp.

Close to *dubitans*, Walk., in appearance. Thorax and fore wings of the same blackish brown as in the darkest *dubitans*, but more smooth and shining and the wings somewhat narrower; a light brown area at centre of inner margin. Lines lost; t.-p. line indicated by fine white venular dots, s.-t. by a white powdering preceded by black, which forms a rather distinct spot at anal angle. Fringe crenulate, containing white specks at the ends of the veins. Orbicular a linear black ellipse, broken above; claviform similarly distinctly outlined in black, the filling as dark as the ground colour; reniform black edged except superiorly, with a double curved white mark outwardly. A few white specks on costal edge. Hind wings gray, irregularly shaded. Fringe, abdomen and whole under surface suffused with pinkish purple. Expanse 36 to 40 mm.

Described from 9 specimens, Kaslo, B. C.

U. S. National Museum, type No. 7326.

Homohadena fifia, n. var.

Differs from badistriga, Grote, in being darker gray, the ground