

another between ordinary spots, a third just below that and a fourth on inner margin. S. t. line pale, irregularly incised opposite cell and at lower third of wing, here cutting through quite to t. p. line; the space between it and t. p. line is dark brownish-black, and, as just mentioned, is cut into three patches by the s. t. line. Beyond s. t. line the wing is pale grayish. Fringe gray and white checkered, with well-marked line at base, which is evenly and neatly cut into short bars by the same white dashes which cut the fringe. Orbicular minute white point. Reniform white-ringed, darker centered, rather narrow. The markings on the fore wing are neat and distinct, giving a well-marked checkered appearance, quite different from any other species in the genus. Hind wing dark fuscous, with a very faint discal dot and mesial band. Fringe pale, with darker mesial band.

Head and thorax concolorous with fore wing, abdomen with hind.

Beneath, fore wings fuscous, with obscure discal mark. Hind wing paler, with distinct discal dot and well-marked mesial band.

Type.—One ♂. Redington, Ariz.

*Oligia ensina*. n. sp.—♂. Expanse, 28 mm.

Fore wing reddish-brown. Basal half-line distinct, double, pale-filled. T. a. double, dark brownish-black, transverse across cell, then somewhat outwardly to inner margin, slightly scalloped. The upper half of wing, between basal half-line and t. a., dark blackish-brown, forming a strongly contrasting subquadangular patch. Median shade not well marked, somewhat irregular and dentate. T. p. line double, pale-filled, inner portion more prominent, slightly scalloped. The space between t. a. and t. p. line is quite evenly coloured, there being, however, some blackish scales along costa, above reniform and a blackish streak beyond it. S. t. line fragmentary, composed of pale blotches between veins, the space between it and t. p. line is slightly darkened, with a well-marked black dash across it below costa, and another beyond lower portion of cell. The terminal space is somewhat paler than subterminal, the veins, however, being rather broadly darkened. A rather faint dark terminal line. Fringe concolorous, somewhat paler at base. Orbicular rather small, concolorous, pale-ringed. Reniform quite strongly inwardly oblique, more or less well-developed tooth projecting inward from lower portion, an outer pale ring, within which is a darker ring, the centre again becoming pale. Lower portion with some blackish scales. Hind wing pale blackish-fuscous, darker outwardly, with a well-marked discal dot. Fringe slightly paler, with slightly darker mesial band. Head and lower half of collar