Base of pleura on sternal side provided with a small chitinized tooth, minutely denticulate; 2nd gonapophyses long, the tips long and widely separated (Japan).....incongruens, sp. n.

Erioptera (Acyphona) incongruens, sp. n.

Small species; light brown, with narrow dark brown pleural stripes; wings thickly spotted with brown.

Male.-Length, 5 mm.

Male.—Rostrum and palpi brown. Antennæ long, segment one brownish-yellow; segments two to eight light yellow; remainder with increasing amounts of brown at their tips, the apical segments all brownish. Front, vertex and occiput dark brown.

Thoracic pronotum brownish-yellow, brown on the sides. Præscutum reddish-brown, with a double median brown stripe; humeral region brighter yellow; sides of the sclerite darkened; scutum, scutellum and postnotum brown. Pleuræ reddish-brown with narrow dark-brown lines, the most dorsal one continuing from behind the fore coxa underneath the wing to the postnotum; the second beginning on the mesosternum running above the middle coxa, becoming very narrow and indistinct before the root of the halter; the last stripe on the metasternum over the hind coxa. Halteres light yellow. Legs: coxa brown; trochanters brownish-yellow. (The legs are all detached and loose in the vials; most of these have the femora largely brown, basal third mostly paler, yellowish; a post median yellow ring, tip usually pale; tibiæ and