

one-half the length of the subcylindric basal enlargement, which latter has a length about three-fourths greater than its diameter, sub-basal and apical circumfili and subapically a moderately thick whorl of long, moderately stout setæ; terminal segment slightly produced, narrowly conical, with a length about three times its diameter. Palpi: first segment short, subquadrate, the second with a length three times its width, moderately broad, the third one-half longer than the second, more slender, and the fourth one-half longer than the third, more slender, the distal three sparsely haired. Eyes large, holoptic. Mesonotum pale yellowish brown, the submedian lines bright yellowish. Scutellum dark brown, postscutellum yellowish brown. Abdomen yellowish transparent, the genitalia relatively very large and subdorsal. Wings hyaline, subcosta uniting with the narrowly scaled costa near the basal third, the third vein a little before the apex; the fifth vein, indistinct distally, unites with the posterior margin near the distal third, its branch near the basal half; halteres whitish transparent. Coxæ, femora, tibiæ and the two basal tarsal segments mostly whitish or whitish transparent, the three distal tarsal segments dark brown; claws long, slender, moderately curved, unidentate, the pulvilli about three-fourths the length of the claws. Genitalia: basal clasp segment short, stout, curved, the sides nearly parallel, the apex roundly emarginate, the distal angles produced, the internal being longer and broader; dorsal and ventral plates indistinct, the harpes produced as two long, slender, chitinized, somewhat sickle-shaped pieces having a length greater than the entire hypopygium and extending posteriorly somewhat like a pair of scissors.

*Female*.—Length 1 mm. Antennæ extending to the base of the abdomen, rather thickly haired, yellowish brown; probably 12 segments, the fifth sessile, broadly subconical, with a length about one-half greater than its diameter; low circumfili at the basal third and apically, a sparse basal and a scattering subapical whorl of long setæ; terminal segment reduced, broadly conical and tapering to a broadly rounded apex. Palpi: first segment with a length three times its diameter, the second a little longer, broader, the third twice the length of the second, more slender and the fourth one-fourth longer than the third, all sparsely haired. Eyes holoptic,