

Calidris arenaria (*Linn.*) SANDERLING.

Fairly common on the sand beaches during the autumn migration, from the middle of September till November.

Totanus melanoleucus (*Gmel.*) GREATER YELLOW LEGS.

Fairly common during the migrations. They pass here going north about the first of May and going south from the middle of July till November. About the middle of last June one was observed here.

Totanus flavipes (*Gmel.*) LESSER YELLOW-LEGS.

Not quite so common as the preceding, and are not seen so late in the fall.

Totanus solitarius (*Wils.*) SOLITARY SANDPIPER.

Quite common from the first of August till October. They frequent small ponds and puddles with soft muddy borders.

Bartramia longicauda (*Bechst.*) BARTRAMIAN SANDPIPER.

Rare. Several were observed on the Grand Pre, September, 1896.

Actitis macularia (*Linn.*) SPOTTED SANDPIPER.

Abundant from the middle of May till September. Frequent the shores of rapid streams and upland ponds.

Numenius hudsonicus (*Lath.*) HUDSONION CURLEW.

Rare fall migrant. Observed in early September.

Charadrius squatarola (*Linn.*) BLACK-BREASTED PLOVER.

A fairly common autumn migrant. August till November.

Charadrius dominicus (*Mull.*) GOLDEN PLOVER.

A few years ago this bird was very common during the fall migration. But the last few years they seem to have decreased rapidly. Not more than one is observed to-day, for every 20 that could be seen six or eight years ago. They frequent the salt marsh, sand beach and upland pasture.

Ægialitis semipalmata (*Bonap.*) SEMI-PALMATED PLOVER.

Quite common on the sand beaches from the middle of July till October. Do not nest here.