

about 19, considerable nervous trouble was inaugurated by grieving over a great bereavement, and the menses became more and more painful. The anguish became such a horror to her that she frequently resorted to morphine, partly to allay pain and partly to procure sleep. Fortunately she had not, as yet, contracted the habit, but the tendency was undoubtedly in that direction. When first consulted by her, examination was not granted. Menses appearing shortly afterward, was called upon to afford relief. Flow was very scanty and clotted. There were sleeplessness, terrific headache, pain in back, constipation, etc. Ergoapiol (Smith) was administered, one capsule every three hours. Flow was considerably increased, there was a gradual lessening of all the suffering, and almost complete relief in twelve hours. This young woman had been placed upon Ergoapiol (Smith), one capsule twice daily for one week preceding appearance of menses, and has passed through several periods with very little suffering. An examination made recently showed a marked retroversion and very sensitive cervix. A properly applied supporter will doubtless work considerable benefit in her case, but it cannot be disputed that the comparatively easy menstruations occurring recently, in spite of the displacement, were due entirely to Ergoapiol.

CASE 4.—Miss ———, aged 18, had always been regular in menstruating. Could get no history of any previous disorder within patient's knowledge. Contracted a heavy cold about time of menstrual epoch, and was much alarmed by non-appearance of flow. Discomfort was not marked. Ergoapiol (Smith), one capsule three times a day, was prescribed. Reported later that flow was established in twenty-four hours after treatment was commenced. The delay in this case was about four days.

CASE 5.—Mrs. ——— consulted me, giving the following history: Three months previously had had a profuse uterine hemorrhage occurring about the time of menstrual period. As she had for a number of years menstruated only at intervals of about six or seven weeks, the fact that menstruation had been suspended for six weeks before the date of trouble was not especially significant. The hemorrhage, which was at no time alarming, had continued for several days. Since that time there had been an almost constant wasting and at times a considerable flow. Her condition was practically invalid. Examination revealed a gaping os, a cervix exceedingly tender and abraded, and a large uterus. Before resorting to curettement it seemed advisable to try other measures. Ergoapiol (Smith), one capsule every three hours, was prescribed. In about twenty-four hours there was a decided increase in the discharge, which consisted of clots and considerable debris. There were some pains, of a cramp-like nature. The discharge began to