

## LIST OF TABLES (continued)

<u>Table Number</u>		<u>Page Number</u>
3-19	Avian and Mammalian Species most likely to be Influenced by a Reduction in Food Resources Due to Acid Precipitation, Feeding Habitat During the Breeding Season and Alternate Food Resources.	3-105
3-20	Comparison of Acid Input and Consumption for Harp Lake Subwatersheds.	3-111
3-21(a)	Ratios of Hydrogen Ion (H) to Nitrate Ion (N) Expressed in Equivalents per Liter for Six Streams with Variable pH in the Dorset, Ontario Study Area for 1976, 1977 and 1978.	3-115
3-21(b)	Ratios of Hydrogen Ion (H) to Sulphate Ion (S) Expressed in Equivalents per Liter for Four Streams with Variable pH in the Dorset, Ontario Study Area for 1976, 1977 and 1978.	3-116
3-22	Periodic pH Depressions Observed in Streams and Lakes with Different Sulphate Loadings.	3-128
3-23	Observed Sulphate Loadings, and Loading Values Predicted by Various Methods, Required to Protect Aquatic Ecosystems.	3-131
3-24	Illustrative Dose/Response/Time Relationships Under Selected Loading Patterns.	3-133 & 3-134
4-1	Effects of Long-Term Controlled Ozone Exposures on Growth, Yield, and Foliar Injury to Selected Plants.	4-5 & 4-6
4-2	Exposure-hours to Ozone at Rural Locations in Southern Ontario for 1980.	4-7
4-3	Ozone Concentrations at Some Locations in New York State.	4-8
4-4	Representative Tolerance Limits to Simulated Acid Precipitation.	4-10
4-5	Acidity Related Reactions Influencing Availability of Several Elements.	4-17