

- mixing with the ovula, 508; opinion of Professor Amici that they come into contact with the ovulum, 508-9; traced into the aperture of the ovulum, 540; the author not so far advanced, 509; origin of the pollen tubes, 509; their growth, 509, 510; a vital action, which may be excited by the stigma of another species, 509-510; their nutriment, whence derived, 510; effect of action of pollen on stigma, 510; subsequent changes in the ovulum, *ibid.*; formation of the embryo, *ibid.*; cellular thread passing from its apex to the apex of the original nucleus, 510, 511; changes in the embryo, 511; seed and its funiculus entirely without vessels, 511; areola, or nucleus of the cell, its position, &c. 511-514; spirally striated cells in, 514, 515, 548; raphides in, 515; mucous tubes in various species, *ibid.*; structure of fibrillæ, constituting the pubescence of the aerial roots in, 548
- Orchidæ and Asclepiadæ, analogy between the two orders, 530; in the presence of an additional part, consisting of a modification or production of the stigma, 530-1; in the frequency and necessity of the agency of insects in fecundation, 531; penetration of pollen tubes into the cavity of the ovarium, 531-2; function of the granules within the pollen tubes in reference to impregnation, 532
- Orchidæ and Asclepiadæ, supplementary observations on the fecundation of, 545-551
- Orchis fusca*, cellular thread of ovulum in, 547
- Orchis Morio*, mucous tubes entering the aperture of ovulum in, 540; cellular thread of ovulum in, 547
- Orchis ustulata*, cellular thread of ovulum in, 547
- Orobanchæ, Mr. Lindley's idea of the placentation of, controverted, 562
- Oudney, Dr. Walter, plants collected by in Central Africa, 257, 259, 260, 272
- Oudneya, characters of the genus, and observations on its structure, 272
- Ovarium, vascularity of, compared with that of the leaf, 379 *note*; theoretical view of the origin or formation of, 555; exceptions to the ordinary condition of, 555-6; origin of various forms of compound, 457-563; supposed exceptions in Orchidæ and Orobanchæ considered, 557-562
- Ovulum, structure of the unimpregnated, in *Cephalotus*, indicates the position of the radicle of the future embryo, 77, 440; structure and development of in *Rafflesia*, and in Phænogamous plants generally, 406-7; in *Hydnora* and *Cytinus*, 407; structure of the unimpregnated, 439-452; in *Kingia*, 439; in *Cephalotus*, 77, 440; generally consists of two coats enclosing a nucleus, 440; origin and relation of the coats, and of the future embryo, 440-1; structure of ovulum, as described by Grew, Malpighi, Camerarius, Morland, Etienne Francois Geoffroy, Claude Joseph Geoffroy, Vaillant, Needham, Adanson, Spallanzani, Gærtner, Turpin, Richard, Mirbel, Auguste de Saint-Hilaire, Ludolf Christian Treviranus, Dutrochet, and Link, 442-7; in what respects the account given differs from all those observers, 447; exceptions to the general structure in *Compositæ* and *Lemna*, 448; importance of distinguishing between the coats of the ovulum and those of the ripe seed, 449; how determinable, 449-452; structure and development of unimpregnated, in Orchidæ, 504-5; in Asclepias, 549-550; the ovulum naked in *Cycadæ*, *Coniferæ*, *Ephedra*, and *Gnetum*, 453; arguments in favour of this view, 454; objections to it, 454-5; origin of the ovulum in *Cycadæ* and *Coniferæ*, 458-9
- Ovules, their marginal insertion in the simple pistilla, and in the component parts of the compound ovarium, shown in various instances, 379 *note*; cases of apparent ex-