Mead, C. V. Webster, O. te list of the

essrs. C. F.

ce to what is Cambridge. , Dr. H. A. r in advance e it a partial nation of it, or the apprere and how ecies in their s, etc., it is ration of its resorted to operation of r successors s have been those of the there is not The student ced progress llections and him.

been quite
is painful to
intomologist
seventh in
Dr. Thomas
Hemiptera.
of a large
ices of their

Emma A. stigations in the precedcorrection of omenclature

the ninth in gist, and of

ur scientific

opriation by rations. In nost entirely th a view of the requisite the Governion, and the t the recomies in British which are the hatching

of the young locusts. This plan is believed to be feasible, as the breeding grounds are not co-extensive with the so-called Permanent region, but are limited to the richer valleys, plateaus and river borders within it.

The Commission will also, it is understood, in its forthcoming report, recommend to the Government a scheme for a system of warning and prevention, through the aid of the mounted police patrol of the Dominion Government, and our signal bureau and

military posts.

Having been favoured with a transcript of the subjects to be treated of in the forthcoming Second Report of the Commission, and the assignment of subjects to the respective members of the Commission, I have no hesitancy in giving assurance of a volume of unusual interest and value. It is to be hoped that Congress will not repeat the inexcusable blunder of ordering of it an edition by far too small to supply the demand, or for the accomplishment of a main object in its labourious preparation—the diffusion of the needed information among those to whom it could not fail of proving beneficial.

The Commission is also occupied with investigation of the Hessian-fly and the Chinch-bug—each of which are chargeable with annual injuries to the amount of

several millions of dollars.

The investigation of the natural history and habits of the Cotton-worm, commenced by the Department of Agriculture last year, has by direction of Congress, been transferred to the Entomological Commission. Prof. Riley has been pursuing its study in Southern Texas and in the Gulf States, aided by special assistants, and it is believed that discoveries have recently been made which will reduce the cost of destroying the

larvæ to perhaps a fourth of what it has hitherto been.

Among the special subjects of study which have claimed attention lately, an interesting one has been the pupation of butterflies. Observations made during the past year on the pupation of some of our butterflies have shown us that we have been at fault in accepting the account given of it by Reaumur over a century ago, and received and quoted by subsequent authors. The most interesting operation in the pupation of the suspensi butterflies is the withdrawal of the chrysalis from the larval skin, the casting off of the skin with its attachment by the terminal legs to a button of silk spun for the purpose by the larva, and the attachment and suspension of the chrysalis by its anal spine to the silk button. Reaumur represented it as accomplished by the chrysalis in its extensions and contractions grasping the larval skin between the segments, and by this means raising itself until it regained the button. Recently Mr. Osborne, an English Entomologist, discovered a membrane serving as a suspensory agent in the change to the pupal state, and for the first, questioned the account given by Reaumur. His observations were confirmed by those of Mr. W. H. Edwards, and followed up by additional observations on large numbers of Nymphalide and Danaide, some of which have been presented in the Canadian Entomologist. There seems to be no question of the existence of such a membrane, and that it consists of the portion of the larval skin lining the region of the rectum, caught upon two knobs conveniently placed for the purpose. Prof. Riley, in a communication to Psyche (vol. ii., p. 249) finds other means of chrysalis suspension—the principal one being the shed intestinal canal, and accessory ones, the tracheal vessels of the last pair of spiracles; these Prof. Riley regards as the principal agents in suspension. In opposition to this, Mr. Edwards considers these ligaments as of but little, if any, service, and finds the membrane to furnish all the requisite support. Additional observations are required to reconcile these different views.

The beds of fossil insects recently discovered in the Tertiaries of our western Territories are proving to be wonderfully rich in number of species and condition of preservation. From a single small basin exposed by a railway cut in the vicinity of Green River Station, Union Pacific Railroad, in Wyoming, Mr. S. H. Scudder in Fossil Insects of the Green River Shales (Bull. U. S. Geolog.-Geograp. Surv. Terr., iv., No. 4, pp. 747-776) enumerates eighty species, representing all the orders of the Insecta except Lepidoptera. An idea of the richness of these beds may be obtained from the statement, that a two hours' search was rewarded by the collection of fifty new species. We are glad to learn that Mr. Sudder is engaged upon a general work on our fossil insects, which will form