

short, but nearly as long as those of the male. Legs short and stout, anterior tibiæ with a short stout spine at either side of tip; middle and posterior tibiæ with minute terminal spurs only. The claws of tarsi are toothed in both sexes.

Primaries with 7 to 10 from a stalk in the order 10, 7, 8, 9; 3 to 5 from end of median, 4 nearer to 5 than to 3.

Secondaries 6 and 7 from same point at end of sub-costal; 8 from sub-costal half way to base; 3 to 5 almost equidistant from end of median.

Moeschler, in 1876, Stett. Ent. Zeit., XXXVII., 297, makes this a synonym of *Spilosoma*.

*L. acraea* Drury.

- 1770—Drury, Exot., I., pl. III., f. 2 ( $\textcircled{F}$ ) and 3 ( $\textcircled{M}$ ), *Phalæna*.
- 1793—Fabr., Ent. Syst., III., 1, 451, *Bombyx acria*.
- 1797—Sm. & Abb.\*, Ins. Ga., II., 133, pl. 67, *Phalæna acria*.
- 1816—Hüb., Veizeichniss, 184, *Estigmene*.
- 1820—Hüb., Samml. Auss. Schmett., II., pl. 404, *Estigmene acria*.
- 1823—Harris\*, Mass. Agr. Repos. & Journ., VII., 323, *Arctia*.
- 1833—Harris, Cat. Ins. Mass., 591, *Arctia*.
- 1837—Westw., ed Dru., I., 7, *Spilosoma*.
- 1841—Duncan\*, Nat. Libr., XXXII., 171, pl. 20, ff. 1-3, *Spilosoma*.
- 1860—Clem., Proc. Ac. Nat. Sci., Phil., XII., 53, *Spilosoma*.
- 1862—Morris\*, Synopsis Suppl., 342, *Spilosoma*.
- 1862—Harris\*, Inj. Ins., Flint, ed. 351, f. 169, pl. VI., ff. 9 ( $\textcircled{M}$ ) and 10 ( $\textcircled{F}$ ), *Spilosoma*.
- 1864—Pack., Proc. Ent. Soc., Phil., III., 125, *Leucarctia*.
- 1869—Bdv., Lep. Calif. (Ann. Soc. Ent., Belg., XII.), 78, *Arctia acria*.
- 1873—Stretch\*, Zyg. & Bomb., 99, pl. 4, ff. 1-3, and pl. 10, f. 6, *Leucarctia*.
- 1874—Morrison, Psyche, I., 21, *Leucarctia*.
- 1875—Edw.\*, Proc. Cal. Ac. Sci., V., 368, *Leucarctia*.
- 1883—Weed, Papilio, III., 84, *Leucarctia*.
- 1883—Grt., Papilio, III., 84, *Leucarctia*.
- 1883—Stretch, Papilio, III., 141, *Leucarctia*.
- 1884—Bean\*, CAN. ENT., XVI., 66, *Spilosoma*.
- 1886—Smith, Ent. Amer., II., 79, *Leucarctia*.