- 78. Goes occulata Lec. Two specimens taken on felled Hickory.
- 79. Acanthoderes dicipiens Hald. Not common.
- 80. Leptostylus aculiferus Say. Two specimens; upon Butternut.
- St. Leptostylus parvus Lec. Very rare; place of capture not recorded.
- 82. Leptostylus sexguttatus Say. Leptostylus commixtus Hald. of former list. Rare. Probably from Hickery.
 - 83. Leptostylus perplexus Hald? One specimen; determination doubtful.
- 84. Leptostylus macula Say. Common; June and July; most frequently on Hickory and Butternut, but has been also taken on Maple and Poplar.
 - 85. Leptostylus collaris Hald. One specimen; probably from Hickory.
- 86. Liopus alpha Say. Liopus cinereus Lec. of former list is now included in this species. Common; on Hickory in midsummer.
 - 87. Liopus punctatus Lec. Two specimens; accidental captures.
- SS. Lepturges symmetricus Hald. Lepturges angulatus Lec. of former list. No record of habits.
 - S9. Lepturges signatus Lec. Very rare. No record of habits.
- 90. Lepturges querci Fitch. Lepturges facetus Say of former list belongs to this species. Common; usually on Hickory, one from Willow.
 - 91. Hyperplatys aspersus Say. Common; upon Hickory, June and July.
- 92. Hyperplatys maculatus Hald. Common; with preceding, and is probably only a form of aspersus.
 - 93. Graphisurus fasciatus Deg. Abundant; infests Hickory and Maple.
 - 94. Acanthocinus obsoletus Oliv. Also common; occurs with preceding.
 - 95. Hoplosia nublia Lec. Rare; June, no record of infestations.
- 95. Pogonocherus penicellatus Lec. Several specimens taken on fences and tree boxes in the city; also one upon dead Pine.
 - 97. Pogonocherus mixtus Hald. Not common; no record of infestation.
 - 98. Eupogonius tomentosus Hald. One specimen.
 - 99. Eupogonius vestitus Say. Two specimens; taken also by Mr. Simpson.
 - 100. Eupogonius subarmatus Lec. Rare; on Scented Raspberry and on Ash.
 - 101. Saperda calcarata Say. Rare; a destructive borer in Poplars.
- 102. Saperda mutica Say. Rare; on Willow; one pair June and one pair July.
- 103. Saperda candida Fab. Not common; occurs upon Hawthorn, Shad-bush and wild Plum.
 - 104. Saperda vestita Say. Abundant; a borer in Basswood.
- 105. Saperda discoidea Fab. Common; June to August on fallen and dead Hickory.
- 106. Saperda tridentata Oliv. Abundant; infests principally the Elm, but also the Maple.

૱.

107. Saperda lateralis Fab. Two specimens; accidental captures.