

29.—WHITE-THROATED GOATSUCKER.

Caprimulgus albicollis, *Ind. Orn.* ii. 585. *Gm. Lin.* i. 1030.

L'Ibijau, *Voy. d'Azara*, iv. No. 310.

White-throated Goatsucker, *Gen. Syn.* iv. 596. *Shaw's Zool.* x. 155.

LENGTH twelve inches, breadth nineteen. Bill brown, with a black tip; nostrils rather prominent; plumage rufous brown, dotted with black; upper part of the body the same, but more obscure; on the throat a large triangular white mark, the feathers fringed with dusky; under parts of the body pale brown, crossed with dusky lines; scapulars, and most of the wing coverts, marked with a black band near the end; the tips yellowish buff; second quills spotted with cream-colour on the outer web; the greater dusky black, crossed about the middle with a white bar; tail somewhat cuneiform; the four middle feathers like the back, barred with dusky; the next on each side white; the last but one white on the inner web, and dusky black on the outer; near the base a white spot; the exterior dusky black, but white on the inner web, near the base; legs brown, middle claw very long, and greatly serrated.

Inhabits Cayenne.—In the collection of General Davies. Is also found at all seasons in Paraguay.

30.—BERBICE GOATSUCKER.

LENGTH scarcely nine inches. Bill stout at the base, with a few strong bristles; plumage above not unlike that of the European Species; round the back part of the neck a rufous crescent; on the wing coverts some mottlings of white; on the throat a large spot of white; on the back a mottled mixture of rufous and white; under parts chiefly waved rufous and dusky, towards the vent whitish; on the first three quills a white spot on the outer web, about the middle,