extended over the remainder of the year. They were made altogether without the aid of a gun, the birds being observed by means of field glasses.

In all there were observed here during the year 142 species, of which 33 were noted by all five observers, 39 by four of them, 25 by three, 13 by two and 32 by one only. The result of these observations is the addition of seven new species (two of which await further confirmation) to the Club list, bringing the total number recorded by the Club up to 222. The additions are as follows:—

- 1. Hydrochelidon nigra surinamensis. Black Tern; already noted by Mr. White in Spring Report.
- 2. Buteo lineatus. Red-shouldered Hawk: one shot at the Quarries on the Montreal Road on 24th Sept. by Mr. White.
- 3. Otocoris alpestris praticola. Prairie Horned Lark: this sub-species should have been recorded before, as it is a summer resident, whilst O. alpestris is a winter visitor. The present sub-species is rather common from about the beginning of March. Mr. White questions this.
- 4. Acanthis linaria rostrata. Greater Redpoll: Mr. White reports this species as having been taken here some years since, but for some reason it has never been recorded by the Club.
- 5. Geothlypis agilis. Connecticut Warbler: this species was observed by Messrs. Ballantyne and Lees: by the latter in the Mer Bleue at Eastman's Springs, July 21, and at Clark's Bush July 22, and by both at Dow's Swamps Aug. 5 and Sept. 2. As this species was only identified with the glass it is inserted as doubtful, pending further investigation.
- 6. Cistothorus palustris. Long-billed Marsh Wren: one specimen shot by Mr. Edwards, Taxidermist, on 21st May, and now in the Liuseum of the Geological Survey.
- 7. Turdus Aliciæ. Gray-cheeked Thrush: several reported shot here and in Prince Edward County, Bowerman. We see no reason why this lird should not have been observed here before, but as there is also doubt as to the identification of this species, it is held under query as in the case of No. 5.