

patients themselves connect with that malady, but rather of those insidious symptoms which we, as medical men, who have lived long enough to have seen years of practice, trace back to a syphilitic basis, hereditary or acquired. A case of recurring rheumatism of this nature, a case of recurring ulceration of the fauces, a case of general nervous exhaustion with flying pains in limbs, loss of appetite, general debility, loss of hair, and remaining thickening enlargement in the groin, a sequence of bubo; these have been instances in which the administration of the bromide of quinine, in doses of from two to three grains three times a day, has been more immediately and determinately beneficial than any other treatment I have practiced myself, or seen practiced by my brethren of physic, in such forms of disease.

One great advantage of this preparation seems to me to be, that it allows one to give much larger doses of quinine than are common, and in frequent and continued doses without setting up the symptoms of headache, oppression, and ringing in the ears, which make what has been called *chinechonism*. Thus we may give three grains of bromide of quinine, three times a day, without inconvenience, for several days, if a smaller dose does not suffice.

I have an idea that the bromide of quinine might be administered with advantage in the earlier stages of the contagious diseases, such as small-pox. It would, I think, allay the severe nervous symptoms which usher in these diseases, and so moderate the secondary symptoms that follow in train. Since I began to introduce the bromide into practice, I have not had an opportunity of putting this suggestion to the test, but I have sent some of the preparation to Mr. Marson of the Small-pox Hospital, asking him to give it impartial trial. I have also asked my friend, Dr. Broadbent, to make trial of it, at the Fever Hospital in all cases of acute febrile disorders. The results they obtain I shall hope to communicate in a future number of this journal.

BROMIDE OF MORPHINE.—Is a useful addition to the salts of the alkaloid. It seems to me that a smaller dose of the salt than is effective in the case of the other morphine salts produces as distinct a narcotic influence, and also that the dose may be repeated more frequently without producing those after effects of an opiate which tell against repetition of administration. For