

Angeioma. Angeioma pigmentosum et atrophicum.  
By WILLIAM A. HARDAWAY, M. D., St. Louis.

Angeioma cavernosum. Lymphangeioma.  
By F. A. HERSMAN, M. D., St. Louis.

Tuberculosis cutis. Lupus vulgaris. Scrofuloderma. Lupus erythematosus.  
By JOHN T. BOWEN, M. D., Boston.

Lepra: *a*, Tuberosa; *b*, Maculosa; *c*, Anæsthetica. Dermatitis venenata. Dermatitis medicamentosa.  
By PRINCE A. MORROW, M. D., New York.

Carcinoma. Epithelioma. Rodent ulcer. Sarcoma.  
By JOHN A. FORDYCE, M. D., New York.

Psorospermosis. Paget's disease.  
By M. B. HARTZELL, M. D., Philadelphia.

Hyperesthesia: *a*, Pruritus; *b*, Dermatalgia. Anæsthesia. Urticaria: Urticaria pigmentosa.  
By EDWARD B. BRONSON, M. D., New York.

Mycosis fungoide. Framboesia. Verruga Peruana. Furunculus orientalis (Aleppo boil).  
By W. HENRY BLANC, M. D., Sewanee, Tenn.

Scabies. Sarcoptes scabei communis. Dermodeæ folliculorum. Pediculosis: *a*, Capitis; *b*, Corporis; *c*, Pubis. Leptus autumnalis. Pulex penetrans. Cimex lectularius. Filaria medinensis. Filaria sanguinis hominis. Cysticercus cellulosæ cutis. Echinococcus hydatid. By FREDERICK LEVISEUR, M. D., New York.

\* COMPLETE IN THREE LARGE OCTAVO VOLUMES.

PRICE PER VOLUME,

Cloth, \$6.50; Sheep, \$7.50; Half Morocco, \$8.00.

Volume I now ready. Volumes II and III in press, and will be issued at intervals of about three months apart.

Orders taken for the complete work only.

D. APPLETON & CO., Publishers,  
1, 3, and 5 BOND STREET, NEW YORK.