

RONCUS PACIFICUS, n. sp.

Length, 1.6 mm. Colour, pale whitish; mandibles and claws of palpi reddish; femora of palpi sometimes reddish; dorsal scutæ of abdomen yellowish-brown; legs with a pinkish tinge; young specimens lighter than adult ones. Mandibles large, as in *Clithonius*; cephalothorax somewhat narrower behind than in front, sides slightly curved, one large eye each side near anterior margin; palpi not as long as body; hand with fingers almost as long as cephalothorax plus mandibles; femur about as long as cephalothorax, sides about parallel; tibia almost conical, almost as wide at tip as the hand at base; hand not pedicellate, longer than tibia, sides but little convex; fingers longer than hand, about straight; legs short, hind pairs not much enlarged. Body and appendages with simple hairs.

Washington State. [Kincaid.] A few specimens.

IDEORONCUS, Balzan.

Allied to *Obisium*, but has only one eye each side of cephalothorax and the claw of mandible has a prominent stylet. Species previously known from South America.

IDEORONCUS OBSCURUS, n. sp.

Length, 3.2 mm. Colour, pale whitish; anterior part of cephalothorax with a reddish tinge, mandibles pinkish, palpi red, dorsal scutæ pale yellowish; body soft. The mandibles have a long stylet, which is divided into two parts; the cephalothorax is a little narrower in front than behind; the eyes, hardly perceptible above, are two, one each side, about its diameter from anterior margin; the cephalothorax just in front of eyes is more suddenly narrowed than elsewhere. Palpi shorter than body; trochanters almost twice as long as wide, the internal margin about straight, external margin quite suddenly but not greatly swollen; femur not quite as long as cephalothorax and broader than trochanters, pedicellate, internal margin at first convex then concave, external margin convex, slightly narrower near tip than near base; tibia shorter, pedicellate, external margin evenly convex, internal one convex, more so near base than near tip; hand as long as tibia, short pedicellate, much broader than other joints, neither margin much convex; fingers shorter than hand, slightly curved, stout. Body and appendages with simple hairs. Abdomen long, sides nearly parallel, not much broader than cephalothorax. The outlines of trochanters can be distinctly seen on the femora