

SCIURUS CAROLINENSIS, Gmel.

Grey Squirrel. Black Squirrel.

Sciurus nige., Rich.*Sciurus leucotis*, Gapper.*Sciurus migratorius*, Aud. and Bach.

From western New Brunswick, through southern Quebec and Ontario as far west as the north shores of Lakes Huron and Superior. Especially abundant, however, in the wooded parts of the Western Peninsula of Ontario.

SCIURUS FOSSOR, Peale.

California Grey Squirrel.

Given in Lord's list of Mammals from southern British Columbia.

TAMIAS STRIATUS, Linn.

Striped Squirrel. Chipmunk.

Sciurus (Tamias) lysteri, (Ray) Rich.

Common from the Atlantic coast westward to eastern Manitoba. Not found west of Lake Manitoba.

TAMIAS ASIATICUS, Gmel.Var. *borealis*, Coues.

Northern Chipmunk.

Sciurus (Tamias) quadrivittatus, Rich.*A-kwin-gwis*. Saulteaux.

Ranges from the western side of Hudson's Bay, on the Churchill and Nelson Rivers to the north shore of Lake Superior, westward to the Rocky Mountains and northward on the Mackenzie River to the Arctic Circle.

Var. *quadrivittatus*, Say.

Rocky Mountains, and mountains in British Columbia in the vicinity of the International Boundary Line.

Var. *Townsendi*, Bach.

Coast of British Columbia.