

the Quaternary fossils of Norway, mentions *Brissus fragilis*, a species which I have not seen, as occurring in the glacial clays of that country (which contain fossils resembling those of the Ottawa clays) and also an undetermined species of *Amphidotus*. The present species may be identical with one of these; but I have no description or figures of Sars' specimens. Perhaps there may be better means of reference to them in the Geological Survey library.

In the meantime I can only say that the specimen probably represents a species of *Spatangus* or *Brissus* which lived in the seas of the Canadian Pleistocene, but which, so far as I know, has not yet been found here in a living state. The fact is another instance of the circumstance which I have noted in papers on the Pleistocene fossils that there is in our Pleistocene marine fauna a North-European aspect, as if at that time the indentations of the two sides of the North Atlantic were more nearly alike than at present.

---

## TWO LAND SHELLS NEW TO THE CANADIAN LIST.

---

By REV. GEO. W. TAYLOR, M.A., F.R.S.C. (Nanaimo, B.C.)

I have the pleasure of recording two additions to our list of Western Canadian land shells.

One is *Punctum Clappii*, Pilsbry, a little shell very nearly allied to *Punctum conspectum*, but not quite so large. *P. conspectum* is common near Victoria, but apparently does not occur at Nanaimo, in which district *P. Clappii* seems to replace it. This latter shell is quite plentiful all round Nanaimo and on Gabriel's Island and is commonly found among decaying leaves, being especially partial to the dead fronds of *Aspidium munitum*. It is also common in Oregon and Washington Territory, where the type specimens were collected.

The second species I have to record is new not only to Canada but to Science. Three specimens were found by me, in 1895, under pieces of board near the Nanaimo water works, and from these as types Dr. Pilsbry has described *Pristiloma Taylori*. I