

after, as he was about to leave the Hospital, there came around the seat of the first sore a plentiful crop of small pimples, which shortly formed into soft chancres. These were gradually healing when he ran away, having violated rules. The use of calomel above referred to, consisted in sprinkling the chancre with it. This has been found useful in several obstinate cases of Hunterian chancre, and is recommended instead of the internal use of mercurials.

Margaret B., aged 70, native of Ireland, admitted October 25, 1870—a burn. Caused by falling, probably while intoxicated, upon a heap of burning shavings. There was extensive burning of the skin upon the left side of the face, neck, chest and over the stomach. There was a good deal of prostration, and it was necessary to support her. The burn was treated by the application of linseed oil and carbolic acid, and the exclusion of the air from ulcerated surfaces. The healing did not quickly set in, but finally proceeded, and all ulcers were closed up by 1st December.

Kate C., aged 21, native of Canada, admitted 2nd November, 1870—typhoid fever. A servant, with well marked symptoms of fever, the prognosis unfavorable. Has been ill for a few days. To have a warm bath, and take fever mixture—liq amm acet., 2 oz., spt. eth nic., 1 oz., tinct. hyos., 6 drs, aqua ad., 8 oz., every three hours, unless in a sound sleep or sweating. In twenty-four hours' time the symptoms were much modified. The condition of the patient did not materially change during the following seven days. In the meantime she took freely of beef tea, milk, and occasionally farinaceous food. On the 11th, whiskey, 4 oz in twenty-four hours was ordered. On the whole, the patient took but little stimulant. Convalescent about the first week in December. At the last, as a tonic, she took tinct nux vom., 10 drops three times a-day.

James G., aged 28, native of England, admitted 24th November, 1870—acute rheumatism. Has been ill two days. Parts first affected were feet, then legs, and then the arms. Treated by administration of pot. iodidi, 5 grs. every eight hours. Hydrate chloral, as an anodyne when necessary, grs. from 10 to 30. On the 26th, a black draught, with 40 drops tinct opii was given. The acute symptoms abated, and became much better until Dec 6, when, from careless exposure or the condition of the weather, he was worse. A good deal of swelling and redness of feet and hands, but a strong lotion of plumbi acet. soon