

congress of Bologna; 46-47. The distinction of eozeic and cenezoic serpentines in Italy; their geographical distribution.

IV.—*Geology of the Alps and Apennines*.—§ 48. Theories as to Alpino crystalline rocks; 49. The work of Gastaldi; 50. Sections by Nori and by Gerlach, in the Western Alps; 51-52. Sections by Gastaldi; ancient gneiss, pietro-verdi, newer gneiss, youngest schists; 53. Characters of newer gneiss; 54. Section in the Biellese; 56-57. The pietro-verdi zone in its wider and its narrower sense; 58. The youngest crystalline schists; 59. Great thickness of these Alpine rocks; 60. Von Hauser's sections in the Eastern Alps; 61. Rocks of the Simplon; Primitive schists of Lory; 62-66. The lustrous schists; 67. Rocks of the St. Gothard; 68. Granites and so-called porphyries; 69. Classification of Alpine crystalline rocks in four groups; 70. Their relations to fossiliferous strata; 71-73. Geology of Corsica and Elba; 74. Geology of Sardina; its Cambrian fauna; Ordovician [foot-note]; 75. Change of views as to the age of crystalline rocks; 76-77. The marbles of Carrara; 78. Studies in the Valtelline; 79. The granulitic series of Saxony; 80. Conglomerates in Saxony and Sweden; 81-82. The Montaiban series defined; 83. Geology of Eastern Bavaria; 84. Bojian and Hercynian gneisses; primitive clay-slate series; 85-86. Gastaldi on the continuity of the Alps and Apennines, and the geognosy of serpentines.

V.—*The Serpentine of Italy*.—§ 87. Pre-paleozoic serpentines; 89. Supposed tertiary serpentines; 90. Gastaldi on the sedimentary origin of serpentines; 91-93. The hydroplutonic hypothesis; Issel and Mazzuoli; 94-95. Pellati's criticisms; 96-97. The ophiolite of Monteferrato described; 98. Inversion of adjacent strata; 99. Constitution of the mass; 100-101. Capacci's view of its origin: recomposed serpentines; 102. Difficulties of this view; 103. Proposed explanation; 104. Relations of older and younger strata; Bonney on the igneous origin of the Monteferrato ophiolite; 106. Ophiolites near Sestri Levante, and near Biella; 107. The Staten Island serpentine.

VI.—*The Genesis of Serpentine*.—§ 108. [a] By metasomatism of plutonic masses; conversion of feldspathic rocks to serpentine; 109. Of feldspathic rocks to limestone; of serpentine to limestone; King and Rowney; 110. [b] By metasomatism of neptunian masses; supposed change of limestones to serpentine; to granite, gneiss and petrosilex; 111. [c] By diagenesis of silico-magnesian sediments; 112-113. [d] By alteration of plutonic olivine-rocks; [e] the hydroplutonic hypothesis; 114 [f] By aqueous deposition of magnesian silicates; 115-116. Dieulefait and Lotti on the sedimentary origin of Corsican serpentines; 117-119. Constitution of serpentine-masses; various magnesian silicates and their relations; hydrous and anhydrous species; examples of their simultaneous formation; 119-120. Alteration of olivine to serpentine; 121-122. Geognostical relations of olivine; its presence in limestones; 123-125. Dunite and lherzolite; their indigenous character; olivine-bearing schists.

VII.—*Stratigraphical Relations of Serpentine*.—§ 126. Contradictory opinions; Geikie and Bonney on Ayrshire serpentines as indigenous and as exotic; 127. analogies from coal-seams; 128. Stapf on the serpentines of St. Gothard; their seemingly intrusive character; 129-130. Aqueous origin of these serpentines and their displacements; the structure explained.

---

*Note.*—In printing the above paper, in October, 1883, the author has referred to results made known to him since it was presented in May, 1883; namely, the recent studies of Lotti, (§ 72, 73) and the experiments noted in §110.—[T. S. H.]