

—Prof. Parvin says that all drugs which by their therapeutic action tend to increase the *Arterial tension* will also increase the flow of the milk.

—As a rule, Prof. Keen says, in every four out of five cases of *Fistula in Ano*, the patient will be found to be of a tubercular character.

—*Malarial Fever*, according to Prof. Wilson, will be rarely met with, if at all, in those regions in which the temperature does not rise above 60° F.

—Glycerine or any of the mineral fats, Prof. Wilson says, should not be employed by inunction in the skin in cases of *Scarlet Fever*, but fresh animal fats should be employed in making inunctions.

—Prof. Hare says that a combination of bromide and caffeine will often cure a *Headache* that neither the bromide nor the caffeine alone will relieve.

—Prof. Parvin says that two hours should be allowed to elapse before active measures are instituted toward the forcible removal of a *Retained Placenta*.

—Prof. Parvin thinks that the *Involution of the Sexual Organs* after confinement takes place more rapidly and satisfactorily if the mother nurses her child.

—Prof. Parvin does not favor the administration of ergot during the *Third Stage of Labor* unless the patient be a habitual bleeder, if hemorrhage be present.

—The *Hemorrhage* occurring in cases of laceration of the cervix. Prof. Parvin has found, can generally be controlled by the injection of hot water into the vagina.

—*Inability to Nurse*, Prof. Parvin says, is often due to heredity; brought on by the fact that a number of successive preceding generations did not nurse their offspring.

—Unless during the existence of an epidemic of smallpox, a child who is suffering from a cutaneous disease, or who is otherwise in poor health, should not be *Vaccinated*.

—Fifteen to twenty grains of bismuth and one to two grains of carbolic acid administered every hour or two, Prof. Hare says, will be found to generally stop *Reflex Vomiting*.

—*Fibromatous Tumours*, Prof. Keen says, are neither painful nor tender to the touch; they are also slow in growth. They may, he says, by pressure on adjacent parts, produce pain.

—The eruptions produced by the inoculation of *Vaccine Lymph* will manifest themselves, Prof. Wilson says, twenty-four hours earlier, if the inoculation has been performed by human lymph, than if performed by the bovine.

—Prof. Wilson says that relapses of *Scarlet Fever* are rare, but a secondary attack may occur at some later period, the first attack not conferring an immunity from subsequent attacks.

—*Cannabis indica*, according to Prof. Hare, will be found to be a very useful drug in stopping the *Cough of Phthisis*, and it possesses the advantage over opium in that it is not so depressant to the system in general.

—*Sarcoma*, Prof. Keen says, as a rule, is a disease of youth and not of old age. It makes its appearance generally during the period when the tissues are growing. It generally appears between the age of twenty and thirty, more so than after forty.

—Prof. Parvin says that during the *Period of Menstruation* the condition of the mother's milk is altered, and often is the cause of an attack of colic in the nursing child. But as soon as the menstrual period has elapsed then the milk returns to its normal condition.

—Camphoric acid, according to Prof. Hare, is the best drug that can be used in controlling the *Nightsweats of Phthisis*. It should be taken in doses of twenty to thirty grains, and two or three hours before the time that the sweats generally come on.

—*Anteflexion of the Uterus*, according to Prof. Montgomery, is of most frequent occurrence in women who have never borne children. It is also the displacement which is found of most frequent occurrence in the sterile woman.

—The tumour which is *Scrofulous* in character, Prof. Keen says, in the early stages will be found to be perfectly movable, but in the later stages it will be bound down tight and will be immovable, due to its having infiltrated into the surrounding tissues.

—The oxalate of cerium, either alone or combined with bismuth, Prof. Hare says, will very often be found to stop *Excessive vomiting*. Especially has this been found so in such cases as are due to a hyperacidity, or to an irritation of the mucous membrane of the stomach.

—Prof. Wilson favors the treatment of *Scarlet Fever* by chloral. Such doses, he thinks, should be administered as to keep the patient under its hypnotic influence to such an extent as to require wakening at the time when food or medicine is to be administered.

—In cases of *Chancroids*, which are indolent in healing, Prof. Horwitz recommends the following:—

R.	Ung. hydrarg. nitrat.,	3j	
	Ung. iodoformi,	5ij	
	Ung. zinci oxidi,	3ss	M.

Sig.—Apply locally.