

ing inflammations giving bromides and soothing anodynes with zinc, arsenic and quinine.

Neuralgia, myalgia or rheumatism of the throat has been described, but difficulty is often found in locating the exact spot from which pain emanates. The removal of an enlarged follicle will sometimes cause pain to disappear, but in hysterical or nervous women there may be no apparent cause, and then we must regard it as a local sign of a general condition. Where no local cause calls for treatment you must direct your energies to the constitutional state.

Neurosis of motion is divided into paralyses and spasms. Paralysis may be due to bulbar disease, diphtheria, or may complicate facial paralysis. The central lesions causing this neurosis are bulbar myelitis, hemorrhage, embolism, tumors and meningitis of base. The result is usually a fatal ending. The treatment would be local blood letting, ice bags to nape of neck ; salines and other antiphlogistic measures in the acute form. In the chronic form there is no treatment of any avail. Many things are advised, but no good results have been got. That paralysis following diphtheria and membranous sore throat usually follows fourteen days after convalescence has begun. Must use Galvanism and Faradism every other day until improvement. Strychnine is useful. May have to feed by stomach tube.

The more severe spasms of the pharynx occur in acute uvulitis, œdema of glottis, hydrophobia, and as reflex signs of central tumors. Remove all possible sources of irritation, as intranasal or other growths, disorders, etc. If a simple neurotic state is the cause, hypophosphites, iron and arsenic are indicated. Diet, exercise and freedom from all excitement are essential.