

No. 450? For occasion of the query see Key, p. 267.

No. 466. A pale form is described from California. See Ridgway  
(Am. Nat., viii, 1874, 111).

**466a. RALLUS ELEGANS AUD.,**

*var. OBSOLETUS* Ridg.

**Californian Rail.**

No. 470. A peculiar variety of the black rail is described from the Farallones by Mr. Ridgway (Amer. Nat., viii, 1874, 111).

**470a. PORZANA JAMAICENSIS (GM.) CASS.,**

*var. COTURNICULUS* Bd.

**Pacific Black Rail.**

No. 472. The relationships of this form to the European *G. chloropus* require further investigation. It will probably stand as

**472. GALLINULA CHLOROPUS LATH.,**

*var. GALEATA* (Licht.) Hartl.

No. 479? Compare Key, p. 282.

No. 489. A lighter colored form of dusky duck, with a buff throat, is described from Florida (by Mr. Ridgway, *l. c.*), where it is resident.

**489a. ANAS OBSCURA GM.,**

*var. FULVIGULA* Ridg.

**Floridan Dusky Duck.**

No. 493? The query indicates probability of only varietal distinction from No. 492.

No. 501? The query indicates probability of only varietal distinction from No. 500.

No. 575bis. A specimen of the European white-winged tern, *Hydrochelidon leucoptera*, was taken in Wisconsin, July 5, 1873, by Th. Kumlein, and presented to the Smithsonian by Dr. Brewer. This is the first instance known of its occurrence in this country.

[575bis.] **HYDROCHELIDON LEUCOPTERA (MEIS.) BOIS.**

**White-winged Black Tern.**