

until they are prepared to admit that the power of the heavy Arctic currents passing over the submerged land and carrying with them their burden of ice, is vastly greater as an agent of denudation than either the rivers or glaciers. Nor must we confine this to the Post-pliocene period. Prof. Hall has shewn that the whole of the vast thickness of the Palæozoic rocks of the Appalachians may be attributed to the carrying power of the same currents which are now piling up banks of Arctic sand and stones along the American coast. Nay more, the history of the land of the Northern Hemisphere throughout geological time has been that of a series of elevations and depressions or gigantic pulsations of the earth's crust, so regular that we cannot hesitate in referring them to some constantly operating law. Every elevation exposed the land to sub-aerial disintegration. Every subsidence scraped and peeled it by the action of the Arctic currents, and thus the carriage of material and the growth of the continents have ever been to the south-west. I cannot leave this subject without according to Dr. Carpenter much credit for contending as he has done for the reality, power, and true causes of these great sub-oceanic rivers, which have played and are playing so important parts as geological agents, that without them it is impossible to account either for the Palæozoic deposits or the Post-pliocene deposits of our North American continent.

But it is time to turn to the second topic which I have marked out for myself in this discourse. In the past summer three lines of geological reconnaissance have been pushed out from the Laurentian and Huronian country of Lake Superior over the plains of Manitoba. One of these, under Mr. Selwyn, followed the line of the North Saskatchewan. The second was that of Prof. Bell on the south branch of the same river and its tributaries. The third was that of Mr. G. M. Dawson on the 49th parallel. All of these have been brought under the notice of this Society in the course of the winter. This great western plain presents first a wide expanse of Cretaceous rocks, apparently not highly fossiliferous and not well exposed, but containing some limestone layers rich in Foraminifera and Coccoliths precisely similar to those of the English chalk. Some of these have been described by Mr. Dawson in our Journal. This is succeeded by vast estuarine and lacustrine deposits of clay and sand, holding brackish-water and fresh-water shells, and beds of lignite with abundant plant remains. The general geological