

Wings—Fore wings sometimes hyaline; costa not brown.

Described from a type and five paratypes, the female in the Peabody Museum referred to above being selected as the type. Two of the paratypes are in the collection of the Boston Society of Natural History, two in the collection of the American Entomological Society at Philadelphia, and the fifth is in the collection of the Conn. Agricultural Experiment Station at New Haven, Conn.

Male—The male differs from the female as follows: Greater part of lateral face of pronotum straw colour; an approximately right-angled band on the mesoepisternum, posterior mesal half of pectus, and basal half of venter entirely, straw colour; intermediate coxæ black only at base above, the posterior coxæ except inside and inner longitudinal half beneath, and the apical two-thirds of the posterior tibiæ black (continued to base above).

Length—Female 11–13 mm.; male 11 mm.

The male is here for the first time described, and is the only one that I have seen. It belongs in the collection of the Boston Society of Natural History.

This subspecies approaches *fisheri* Rohwer from Maryland, and may prove to be the same.

Habitat—Sharon, Auburndale, and Woods Hole, Mass. (June and July); Farmington (Norton) and Westville, Conn. (June). I have records also from New York, New Jersey, and Virginia, but they may refer to *lobata lobata*.

INSECT IMPORTATIONS INTO NEW JERSEY DURING THE SPRING OF 1915.

BY HARRY B. WEISS, NEW BRUNSWICK, N.J.

The following insects arrived in New Jersey during the spring of 1915 on nursery stock imported from various countries in Europe. Practically all were alive when taken and many came over in numbers sufficient for them to gain quite a foothold. Identifications, for the most part, were made by specialists through the courtesy of the U. S. Bureau of Entomology.

October, 1915.