

below with mostly stellate spreading hairs, glabrous above or but sparingly puberulent, a foot high; leaves small and narrow,  $\frac{1}{2}$  inch long or less, the lower very rarely few-toothed, the cauline sagittate at base; flowers very small, pale rose-colour, 2 lines long; pods very narrow, 1 to  $1\frac{1}{2}$  inches long and about  $\frac{1}{2}$  line broad, glabrous, slightly curved, mostly divaricate on very slender pedicels 2 to 4 lines long, acute, the stigma nearly sessile; seeds (immature) approximately 1 rowed, apparently wingless; near *A. hirsuta*.

Gravelly banks, Revelstoke, B.C., May 13th, 1890. (*John Macoun*.)

*TRIFOLIUM PROCUMBENS*, Linn.

An erect form of this plant was found by Prof. Macoun in 1893, growing in fields at Comox, Vancouver Island. Not recorded before from western Canada, though the var. *minus* is common on Vancouver Island.

*TRIFOLIUM INVOLUCRATUM*, Willd.

Collected at Revelstoke, B.C. in 1890 by Prof. Macoun. Abundant on Vancouver Island, but not before collected in the interior of British Columbia.

*TRIFOLIUM MICROCEPHALUM*, Pursh.

Collected at Revelstoke and Sproat on the Columbia River, B.C. in 1890 by Prof. Macoun. Common on Vancouver Island, but not before recorded from interior of British Columbia.

*LOTUS CORNICULATUS*, Linn.

Recorded from New Brunswick. Collected in 1890 at Victoria, Vancouver Island, by Rev. Edw. L. Greene.

*ASTRALAGUS ROBBINSII*, Gr. var. *occidentalis*, Wat.

Not before separated in Canada from *A. alpinus*, the western form of which it somewhat resembles. Bow River at Morley, Alberta; near the Glacier at Lake Louise, Rocky Mts.; Deer Park, Columbia River, B.C. (*John Macoun*.) Gui-