

Mrs. C., referred by Dr. F. A. Clarkson, small rodent ulcer just below the eye, which quickly responded to treatment.

Mrs. M., referred by Dr. Barber, of Burk's Falls, rodent ulcer of temporal region. Radium was used on two separate occasions, and the ulcer is now quite healed.

*Epithelioma of the Lip.*—Two cases of superficial epitheliomata of the lip have responded splendidly to treatment. Other epitheliomata have been referred on which prolonged treatment will have to be carried out, and on which we hope to report more fully later. To mention a few:

Mrs. B., referred by Dr. S. M. Hay, for an epithelioma of the buccal mucous membrane, which had recurred after removal. She was seen six weeks after treatment, and there had been no re-appearance; she will, however, have to be kept under observation from time to time.

A case of epithelioma of the lip, with glandular involvement, considered inoperable, referred by Dr. Howitt, Guelph.

Mr. B., referred by Dr. Partridge, of Burk's Falls, with a recurring epithelioma of the buccal mucous membrane.

Mr. A., referred by Dr. N. A. Powell. It is an epithelioma of the lip in which extensive dissection had been done. There was constant pain and two masses which had recently begun to increase in size. Since beginning treatment the pain has ceased, but it is too early to report further as yet.

*Fungating Epitheliomata.*—Fungating cutaneous epitheliomata are particularly suited for radium action, and various techniques can be adopted depending on the individual case under observation. "Cross-fire" action often gives excellent results with the use of different forms of filters. A preliminary curettage and removal of the vegetations is of help in decreasing the time required for cure, but is not absolutely necessary.

T. F. T., aet. 54, referred by Dr. Bowman, of Penetanguishene, Ont., presented on Oct. 29th a fungating mass, as large as a fifty cent piece, below and behind the left ear. There had been a small ulcer for about five years, but latterly the growth had been very rapid. The growth was covered with cauliflower excrescences, and projected  $\frac{3}{4}$  of an inch above the surrounding skin. The edges were hard and everted, and the tissues about were quite hard, as though the growth extended to some depth. There were no enlarged glands to be felt. Under local anesthetic the vegetations were removed, and the next day radium applications were made. These were repeated for four days, and then the patient returned home. He was seen again in three weeks, at which time all that was observed