CONTRIBUTIONS TO CANADIAN BOTANY. 1

By James M. Macoun, Assistant Naturalist, Geological Survey of Canada.

ANEMONE HUDSONIANA, Rich.

Frenchman's Bay, near Southampton, Ont. Aug. 28th, 1901. (John Macoun.) Southern limit in Ontario.

AQUILEGIA COCCINEA, Small.

Niagara, Ont.; Câche Lake, Algonquin Park, Ont.; Otterburne, Man.; Brandon, Man. (John Macoun.) Wingham, Ont. (J. A. Morton.) Grindstone Point, Lake Winnipeg. (J. M. Macoun.) Our only specimens of A. Canadensis are from Ottawa, Belleville and Red Rock, Ont. A. coccinea is easily separable from A. Canadensis either in flower or fruit. In flower by its stout spur which is more than twice the length of that of A. Canadensis and abruptly narrowed near the apex. The follicles of A. coccinea are straight and much longer than the spreading follicles of A. Canadensis.

AQUILEGIA VULGARIS, L.

Roadside, Wyoming near Petrolia, Ont. (John Macoun.)

LESQUERELLA NODOSA, Green, Pittonia, vol. IV, p. 309.

On sand, Castellated Rocks, Milk River, Assa., July 13th, 1895. Herb. No. 10,313.2 (John Macoun.)

LESQUERELLA VERSICOLOR, Greene, Pittonia, vol. IV, p. 310.

On rocky slopes, Stony Mt., Man., June 4th, 1896. Herb. No. 12,401. (John Macoun.)

LESQUERELLA MACOUNII, Greene, Pittonia, vol. IV, p. 310.

On prairies at the police barracks, Medicine Hat, Assa., Aug. 9th, 1895. Herb. No. 10,308. (John Macoun.)

¹ Published by permission of the Director of the Geological Survey of Canada.

²These numbers are those under which specimens have been distributed from the Herbarium of the Geological Survey of Canada.