

*Papilio turnus*, Linn.

Common all through the summer.

*Papilio cresphontes*, Cram.

Common from April to September.

*Papilio asterias*, Fabr.

Common through the summer.

*Papilio troilus*, Linn.

From April to October may be seen; not very common at any time.

*Papilio philenor*, Linn.

Common; found from April to middle of October.

*Ancyloxypha numitor*, Fabr.

Not uncommon in the fall.

*Pamphila zabulon*, Bd.-Lec.

Common.

*Pamphila huron*, Edw.

Have taken a few specimens in October.

*Pamphila peckius*, Kirby.

Common in latter part of summer.

*Pamphila cernes*, Edw.

Common; most so of its genus; found all through summer and fall.

*Pamphila verna*, Edw.

Nearly as common as "*cernes*", and found with that species.

*Pyrgus tessellata*, Scud.

Common in October—not very much before.

*Nisoniades juvenalis*, Fabr.

Common on flowers in September; seen occasionally during the summer.

*Pholisora catullus*, Fabr.

Not uncommon; found principally on flowers.

*Eudamus bathyllus*, S. & A.

Rare. I have only found this east of Marshall, along the C. & A. R. R., and only in October.

*Eudamus tityrus*, Fabr.

Common all through summer.

*Paphia troglodgta*, Fabr.

Not uncommon; mostly seen in September, and delights in resting on dead brush, where it may be easily taken.

*Callidryas sennæ*, Linn.

Rare through summer, and very hard to capture; but in latter part of September is common, and while resting on flowers may be taken with the thumb and finger.