	Abdomen short, cylindrical. Black, with bright golden opaque pollinose markings on thorax and abdomen. See "Laparus" pictitarsis Bigot.
28.	Large species; face bare with bristles on the oral margin, in profile straight or concave, the oral margin projecting. 29 Smaller species, face convex on lower portion, not projecting. 30
20.	Elongate; fourth posterior cell closed, usually a considerable distance before the margin Deromyia Less elongate; fourth cell open or closed in the margin, rarely a little petiolate Saropogon
30	Head broad and flat, face not gibbose, third joint of antennæ slender, elongate
	slender, elongate
31.	Abdomen finely punctulate, of nearly equal width; hind tibiæ at the tip and their tarsi only a little thickened; wings hyaline. TARACTIOUS
	Abdomen smooth; hind tibiæ at the tip and their tarsi thick- ened (compare here Nicocles? scitula); antennal style slender, acuminate; wings variegated
32.	Abdomen flattened; in the female a little broader beyond the middle; in the male near the tip, the last two segments of which conspicuously silvery above NICOCLES Abdomen of nearly equal width throughout, the last segment in
	the male not silvery, fourth posterior cell narrowed. BLACODES
33.	Antennæ without a terminal bristle (Laphrinæ)
34.	The veins at the distal ends of the discal and fourth posterior cells in the same straight line, continuous, or nearly so. 35
	The veins, etc., not in the same straight line, the vein closing the fourth posterior cell more or less remote and oblique. 36
35.	First joint of antennæ clongate, slender; second posterior cell narrowed toward the margin of the wing CEROTAINIA
	First joint of the antenna much shorter than the third; first posterior cell narrowed or closed, the second wide open. Atomosia