

The Canadian Entomologist.

Vol. XXXI.

LONDON, OCTOBER, 1899.

No. 10.

TABLES FOR THE DETERMINATION OF THE GENERA OF COCCIDÆ.

BY T. D. A. COCKERELL, N. M. AGR. EXP. STA.

SUBFAMILIES.

- Males with compound eyes. 1.
Males with simple eyes. 3.
1. Anal ring with hairs (♀). *Ortheziinae*.
Anal ring hairless (♀). 2.
2. Mouth-parts present in adult ♀; legs present in all
stages. *Monophlebinae*.
Mouth-parts absent in adult ♀; legs absent in intermediate stage
of ♀. *Margarodinae*.
3. Abdomen of ♀ terminating in a compound segment; anal orifice
hairless. 4.
Abdomen of ♀ not so terminating. 5.
4. Insects with a scale formed entirely of secretory matter without
admixture of the exuviae; adult ♀ retaining legs and
antennæ. *Conchaspinae*.
Insects with a scale composed partly of the exuviae; adult ♀ with-
out legs. *Diaspinae*.
5. Insects enclosed in a resinous cell with three orifices; adult ♀
apodous, with the terminal segments produced into a tail-like organ,
bearing at the extremity the anal orifice; a prominent spinelike
organ above the base of the caudal extension. *Tachardiinae*.
Not so. 6.
6. Females with the posterior extremity cleft; anal orifice closed above
by a pair of triangular plates. *Lecaniinae*.
Not so; triangular anal plates absent. *Coccinae*.
- In preparing the above table, I have borrowed in places from that of
Green, *Coccidæ of Ceylon*, p. 12.