

Also, in the same dose,—

Calcined magnesia,
Sulphur, sublimed,
Cream of tartar, of each, ʒi ;
Essence of anise, ʒiiss .

When the liver is sluggish, in the morning, once or twice a week, give the following in cachet, or in honey, or sugar water :

Calomel,
Scammony, of each, gr. iii . to gr. v .

For a purgative :

Boiling water, fʒiii ;
Manna in tears, fʒi ;
Senna leaves, ʒi ;
Powder of parched coffee, ʒiiss .

Strain, and take during the day.

With atony of the intestines :

Tr. nux vomica, ʒss ;
Tr. belladonna,
Tr. anise,

Tr. cascara, of each, ʒiiss .

Given in water (8 to 12 drops) before meals.

For biliary calculus in infants, in the intervals, bicarbonate of sodium in grain doses for ten days. The following ten days,—

Syrup of ether,
Syrup of turpentine, of each, fʒiii .

A dessertspoonful before meals.

Twice a week add to the above a coffee-spoonful of—

Sulphur, sublimated,
Cream of tartar,
Magnesia, of each, ʒv ;
Essence of anise, ʒxv .
Before meals.

The following table of foods may prove useful :

Foods permitted.—Milk, cream, and fresh cheese; soups, eggs; all meats in small quantities, especially chicken; legumes, well cooked and when green; potatoes; dried fruits, preferably cooked; marmalades and jams; cooked fish; bread in small quantity; alkaline waters.

Foods not permitted.—Butter and fats, old cheese, pork, mushrooms and truffles, pastries and sweetmeats, dried legumes, sausages, asparagus and tomatoes, liquors, coffee, wines, and strong beers.

For Nephritic Colic.—Hot baths, afterwards blisters, hot poultices, or hot fomentations or stupes applied over the renal region; then administer every half-hour a dessertspoonful of the following mixture :

Antipyrin, gr. viii . to gr. xv ;
Chloroform water, fʒi ;
Lime-water, fʒii ;
Syrup of ether,
Syrup of belladonna, of each, ʒiiss ;
Syrup of orange-flowers, ʒiiss .

If this is vomited, then give morphine hypodermically.—E. PÉRIER (*Revue Obstet. et Gynécol.*, August, 1894).

THERAPEUTIC BRIEFS.

—FOR PSORIASIS:—

R. Ichthyol,
Acid. salicylic.,
Acid. pyrogalllic.,
Aristol, aa gms. $2\frac{1}{2}$
Vaselin,
Adipis,
Lanolin, aa gms. 30. M.

A powerful ointment, to be used in small quantities.

—FOR CHRONIC CONSTIPATION (*Gazetta Medica di Roma*):—

R. Aloës, gr. iv
Strychniæ sulphat., gr. $\frac{1}{4}$
Extract. belladonnæ, gr. $\frac{1}{4}$
Ipecac. pulv., gr. vss . M.
Divid. in pil. xij .

SIG.—One every evening.

—In the German army the following application is employed for the rapid cure of BLISTERS of the feet incident to long marches (*Therapeutic Gazette*):—

R. Saponis nigri, p. 52
Aquæ, p. 27
Vaselin., p. 15
Zinci oxidi, p. 6
Essentiæ lavandulæ, q. s. M.

—From *Medical Press and Circular* of a recent date we quote the following prescriptions:—

Application for CHRONIC PHARYNGITIS:—

R. Iodi, gr. vj
Potassii iodidi, gr. xij
Mentholi, ʒj .
Glycerini, q. s. ad ʒj . M.

Apply with a camel's-hair brush twice or thrice daily.

Useful in BRONCHITIC ASTHMA:—

R. Potassii iodidi, ʒij
Ammon. carb., ʒj
Tinct. lobeliæ, ʒij
Sp. chloroformi, ʒijv
Vin ipecac., ʒij
Infus. senegæ, q. s. ad ʒʒvj M.

A tablespoonful in a wineglassful of water every four hours.

INCONTINENCE OF URINE:—

R. Tincturæ belladonnæ,
Tincturæ cubebæ, aa ʒij
Tincturæ nucis vomicæ,
Tincturæ rhei aromaticæ,
of each, ʒj
Tincturæ cascarillæ, ʒij

12 drops at bed-time for a child from seven to ten years.

The REMOVAL OF WARTS:—

R. Hydrarg. bichlor., gr. v
Acid. salicyl., ʒj
Collodii, ʒj M.