

quent in Middlesex in heavily timbered woods. In Kent, breeds abundantly.

2. *Hylocichla fuscescens*; Wilson's Thrush. Breeds abundantly.

3. *Hylocichla ustulata swainsoni*; Olive-backed Thrush. Very rare in migrations near London; only one shot; more are found in Kent where it may breed; several seen May 19th, 1882.

4. *Hylocichla ustulata palmeri*; Hermit Thrush. Regular migrant. Found common in full song in a large swamp, June 22nd, 1882. No nests found, although it was undoubtedly breeding. None observed in summer in any other locality.

5. *Merula migratoria*; Robin. Breeds abundantly. In very mild winter a few remain.

6. *Mimus polyglottus*; Mock bird. One taken in June, 1860 at Chatham. (E. W. Sandys).

7. *Geothlypis trichas*; Catbird. Breeds abundantly. Five eggs not at uncommon set.

8. *Harporhynchus rufus*; Thrasher. Much less abundant than the last. Seems to be very locally distributed, frequenting low scrub and hazel thickets; breeding in brush heaps, bushes and on the ground impartially.

9. *Sialia sialis*; Bluebird. Breeds abundantly. Earliest eggs, 11th April, 1877; largest set, seven; four sets of white eggs taken.

10. *Poliophtila caerulea*; Blue-gray Gnatcatcher. Generally common, and well distributed through high woods, but some seasons rare.

11. *Regulus calendula*; Ruby-crowned Wren. Common. Arrives from the north early in October, and in mild winters remains, leaving about May 1st.

12. *Regulus satrapa*; Golden-crowned Wren. More common than the last, with same habits. Male in full song shot in a swamp near Lucknow, Bruce County, May 21st.

13. *Parus atricapillus*; Chickadee. Very common in spring and fall; less so in winter, and a few remain through the summer and breed.

14. *Sitta Carolinensis*; White-bellied Nuthatch. Like the last, only more abundant in summer, and less so in winter. Never wakes holes for itself for breeding purposes.

15. *Sitta Canadensis*; Red-bellied Nuthatch. Rare; our few have been taken early in May.

16. *Certhia familiaris rufa*; Creeper. Common, except in summer when a few breed. Nests are placed behind a large flake of bark on the side of a tree.

17. *Troglodytes aedon*; House Wren. Common. Breeds. Snake skins are a common ingredient of country nests.

18. *Anorthura troglodytes hyemalis*; Winter Wren. Common in spring and fall. A few breed in wooded swamps.

19. *Telmatochlamys palustris*; Long-billed Marsh Wren. Breeds very abundantly in the St. Clair

marshes, but in smaller inland marshes is very rare.

20. *Cistothorus stellaris*; Short-billed Marsh Wren. Rather common in the St. Clair marshes; a set of five eggs, the size and shape of those of this species, white and sparsely dotted with the color of those of the other species, was taken at the St. Clair marshes June, 1881, and probably belong to this species.

21. *Anthus ludovicianus*; Titlark. Abundant at times in fall and spring; seen as late as May 20th, and early in spring, flocks of a hundred have been noted.

22. *Monilla varia*; Black and white Creeper. Common in the deeper woods. Breeds.

23. *Helminthophaga chrysopetra*; Golden-winged Warbler. Regularly distributed and rather common. Breeds. Marked very rare by McIlwraith.

24. *Helminthophaga ruficapilla*; Nashville Warbler. Breeds commonly in swamps mostly evergreen, where there is thick growth of moss on the ground, among which the nest is placed.

25. *Helminthophaga peregrina*; Tennessee Warbler. Common at times during spring migration.

26. *Parula Americana*; Blue yellow-backed Warbler. Rather common. Breeds.

27. *Perisoreus tigrina*; Cape May Warbler. Only one taken; Mitchell's Bay, May 16th 1882.

28. *Dendroica aestiva*; Yellow Warbler. Breeds abundantly.

29. *Dendroica cerulea*; Black-throated blue Warbler. Not uncommon during migrations. May breed as a male in full dress was taken June 22nd, 1882.

30. *Dendroica coronata*; Yellow-rump Warbler. Common in spring migrations, more abundant in fall.

31. *Dendroica maculosa*; Black and yellow Warbler. Rare migrant. May breed as a male was taken June 8th, 1882. More common north and east of these counties.

32. *Dendroica cerulea*; Blue Warbler. Common. Breeds. Generally stays high in the trees.

33. *Dendroica Pennsylvanica*; Chestnut-sided Warbler. Common. Breeds.

34. *Dendroica castanea*; Bay-breasted Warbler. Very rare during the migrations. Four specimens only taken.

35. *Dendroica striata*; Black-poll Warbler. Common some seasons during migrations; rare at others.

36. *Dendroica Blackburnian*; Blackburnian Warbler. Somewhat common. Breeds. Frequents the high trees in swampy districts. Often feeds on the ground at Mitchell's Bay.

37. *Dendroica virens*; Black-throated Green