

- Lunatia heros.* Dead specimens occur at North Summerside and Cocagne. Frequently they are grown to the oysters.
- Turbonilla interrupta.* More or less common at Cocagne and at Summerside.
- Odostomia trifida.* Abundant at Cocagne and Bedeque.
- Odostomia fusca.* Rare at Bedeque.
- Odostomia seminuda.* Abundant at Summerside.
- Cerithiopsis Greenii.* Not infrequent at Summerside. Occurs also at Bedeque.
- Bittium nigrum.* Common at Bedeque and occasionally found at Summerside.
- Littorina palliata.* Rare at Summerside, *i. e.*, in the oyster beds.
- Littorina tenebrosa.* One or two small specimens found at Summerside.
- Lacuna neritoidea.* One specimen (probably of this species) at Cocagne.
- Rissoa minuta.* Occasionally at Cocagne. Specimens not extra fine.
- Crepidula plana.* Very common and fine at Summerside. Occurs also at Cocagne and Bedeque. An interesting specimen in its variations, broad, long, flat, curved, and other shapes to suit its position. It lives attached to the shell of the oyster. Color varying from pure white to pea green. Largest about one inch long.
- Crepidula fornicata.* By far the most common, occurs in thousands, from one and a half inches in length to the minute young forms. It also clings to the oyster shell, or to those of its own species, sometimes there being three piled one on another, the lowest clinging to the oyster. Color externally is usually a dingy green or light cream mottled with brown. The interiors of no two are alike, striped, mottled, etc., with shades of maroon and cream color. One specimen had an interior almost jet black, and another is nearly pure white.
- Utricularia canaliculatus.* Occasionally at Summerside. One specimen from Cocagne shows that more might be found by searching.
- Anomia ephippium.* Occasionally at Summerside.
- Ostrea Virginiana.* Has been reported from all these places, but as it is the companion of all in this list, I include it.
- Mytilus edulis.* Abundant at Cocagne, occasional at Summerside, occurs also at Bedeque. This species presents interesting variations. On the south shore of the Province it is narrow with deeply arched valves, the interior nearly black in color. At Bedeque and Summerside the shape is broad and flat with a conspicuous white lining. There is a marked difference between the two places, however; a difference which looks like quality, Summerside shells being much finer in color and smoothness, those at Cocagne having a dingy look. The variety *pellucidus* occurs, and yields very fine specimens. A striped variety also occurs, having parallel bands of black and amber running lengthwise.

Cardium p
Venus me
 Summ
 anoth
 some
 to ha
 white
 Agai
Tellina ter
 but i
Petricola
Saxicava n
Pandora t
Cumingia
 being
Corbula c
Mastra la
 St. Steph