

*restris* the abdomen is black, with a yellow band, and the two last anal segments are white; there are no yellow hairs on the metathorax, and the wings are much clearer.

380. *BOMBUS DERHAMELLUS Kirby*.—Length of body 8 lines. Taken with the preceding.

♀. Body hairy, black. Head with a tuft of yellowish hairs on the vertex; thorax yellow, black between the wings; wings more embrowned than in the male; abdomen yellow at the base with a black posterior band; anus ferruginous.

[274] 381. *BOMBUS PRATICOLA Kirby*.—Length of body 7 lines. Taken with the preceding.

♀. Body black, clothed above with yellowish hairs. Head with a tuft of yellowish hairs below the antennæ, and on the vertex; thorax black between the wings, which are embrowned; legs with yellow hairs at the base; anterior half of the abdomen yellow, posterior ferruginous.

382. *BOMBUS VIRGINICUS Linn.*—Length of body  $8\frac{1}{2}$  lines. Locality uncertain.

♀. Hairs of the body in general black, except a tuft on the vertex behind the antennæ, the anterior and posterior extremities, and sides of the thorax, and the first segment of the abdomen, which are clothed with yellowish hairs; between the wings the thorax is black; the tarsi are rufous; the wings are rather embrowned, most so at the apex; nervures black.

---

#### ENTOMOLOGICAL CLUB OF THE AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE.

The Annual Meeting of the above Club will be held in St. Louis, Mo., on Tuesday, August 20th, the day preceding the meeting of the Association. It is hoped that all Entomologists who can possibly do so will be present on that occasion.