

*Crinus scrophulariæ*, Auct.

A unique of this European *Curculio* was taken on wood wharf, June 30th. There can be little doubt of the correctness of this determination. My specimen is 0.16 inch long (exclusive of rostrum). Body black. Rostrum shorter than thorax, slightly arcuate; antennæ inserted about two thirds from tip, piceous; club somewhat paler. Thorax much narrower than base of elytra, covered with dense prostrate yellowish hairs, and with an elevated transverse ridge at middle; elytra somewhat triangular, blackish, coarsely punctured and with four black longitudinal vittæ, dotted with whitish. A rather large, oblong, dense black spot, margined posteriorly with whitish hairs, placed on the suture, near the scutellum, and a similar but smaller spot at apex. Legs blackish, more or less speckled with whitish hairs.

## ANTICOSTI COLEOPTERA.

COLLECTED ON THE ISLAND IN 1873, BY WM. COUPER, MONTREAL.

(As determined by J. L. Leconte, M. D.)

- |                                                         |                                                         |
|---------------------------------------------------------|---------------------------------------------------------|
| <i>Notiophilus sibiricus</i> , <i>Motsch.</i>           | <i>Chrysobothris trinervia</i> , <i>Lec.</i>            |
| <i>Carabus palustris</i> , <i>Fischer.</i>              | <i>Elatér mixtus</i> , <i>Lec.</i>                      |
| “ <i>lapilayi</i> , <i>Laporte.</i>                     | <i>Agriotes fucosus</i> , <i>Lec.</i>                   |
| <i>Calathus ingratus</i> , <i>Dej.</i>                  | <i>Limonius quercinus</i> , <i>Dej.</i>                 |
| “ <i>confusus</i> , <i>Lec.</i>                         | <i>Sericosomus incongruus</i> , <i>Lec.</i>             |
| <i>Pterostichus coracinus</i> , <i>Lec.</i>             | <i>Corymbites resplendens</i> , <i>Esch.</i>            |
| “ <i>lucotii</i> , <i>Lec.</i>                          | “ <i>aeneicollis</i> ( <i>Kendalli</i> ), <i>Kirby.</i> |
| <i>Harpalus pleuriticus</i> , <i>Kirby.</i>             | “ <i>spinosus</i> , <i>Lec.</i>                         |
| <i>Boletobius pygmaeus</i> , <i>Mann.</i>               | “ <i>aratus</i> , <i>Lec.</i>                           |
| <i>Creophilus villosus</i> , <i>Kirby.</i>              | <i>Eros coccinatus</i> , <i>Say.</i>                    |
| <i>Anthobium dimidiatum</i> , <i>Mels.</i>              | <i>Podabrus basillaris</i> , <i>Say.</i>                |
| <i>Necrophorus vespilloides</i> , <i>Herbst.</i>        | “ <i>laevicollis</i> , <i>Kirby.</i>                    |
| “ <i>mortuorum</i> ( <i>pygmaeus</i> ),                 | <i>Telephorus fraxini</i> , <i>Say.</i>                 |
| <i>Kirby.</i>                                           | <i>Dolichosoma foveicollis</i> , <i>Kirby.</i>          |
| <i>Hydnobius substriatus</i> , <i>Lec.</i>              | <i>Hadrobregmus foveatus</i> , <i>Kirby.</i>            |
| <i>Epuraea boreella</i> , <i>Er.</i>                    | <i>Spondylus upiformis</i> , <i>Mann.</i>               |
| <i>Omosita colon</i> , <i>Er.</i>                       | <i>Criocephalus agrestis</i> , <i>Kirby.</i>            |
| <i>Byrrhus americanus</i> , <i>Lec.</i>                 | <i>Pachyta monticola</i> , <i>Rand.</i>                 |
| <i>Heterocerus substriatus</i> , <i>Kies.</i>           | <i>Acmaeops pratensis</i> ( <i>strigilata</i> .)        |
| <i>Aphodius fimetarius</i> , <i>Ill.</i>                | <i>Leptura pedalis.</i>                                 |
| “ <i>uricola</i> , <i>Mels.</i>                         | <i>Donacia emarginata</i> , <i>Kirby.</i>               |
| “ <i>granarius</i> , var. <i>spretus</i> , <i>Hald.</i> | <i>Syneta tripla</i> , <i>Say.</i>                      |