

DIARRHŒA OF TYPHOID FEVER.

R	Acidi sulphurici aromatici.....	3 ij.
	Ext. hamatoxylin. fld.....	3 ss.
	Spirit. camphoræ.....	3 ss.
	Syrupi zingiberis..... q. s. ad.	3 ij.

M. Sig. Two teaspoonfuls when the stools exceed four in twenty-four hours.—PROF. HARE.

TO RELIEVE THE PAIN IN CYSTITIS.

R	Ext. hyocyami.....	gr. j.
	Camphoræ monobrom.....	gr. ij.
	Morphin sulphat.....	gr. ss.
	Cocoa butter..... q. s.	

M. Et. ft. suppos. No. j. .

WHEN URÆMIA THREATENS.

Use the hot pack over the kidneys, and give:

R	Pilocarpine.....	gr. ss-j.
	Hydrochloric acid dilute.....	3 ij.
	Aquæ dist..... q. s. ad.	3 ij.

M. Sig. Teaspoonful every three hours.—N. A. KREMER, in *The Medical and Surgical Monitor*, Vol. II., No. 6.

CROUP.

The following prescription, suggested by Joseph Holt, of New Orleans, has been tried many times with the happiest results:

R	Chloral.....	gr. lxxv.
	Potassii bromidi.....	gr. xlv.
	Ammonii bromidi.....	gr. xxx.
	Aquæ cinnamomi.....	3 ij.

Sig. Teaspoonful, and repeat in twenty minutes if not relieved.

This is intended for a child about 7 years old. For younger children the dose is slightly diminished. The chloral relieves the spasm of the larynx, and the bromides allay the nervousness; so that the patient is soon asleep, and wakes the next morning as well as usual. The first dose generally gives relief. This prescription is of no benefit in true or membranous croup.—*Practitioner*, 1899, lxiii., 356.

THREADWORMS.

In the treatment of threadworms large injections must be used, and in view of the difficulty of dislodging the worms