

MICRO-LEPIDOPTERA.

BY V. T. CHAMBERS, COVINGTON, KY.

Continued from page 12.

GRACILLARIA.

5. *Gracillaria (Coriscium?) albinatella*. N. sp.

Second joint of the labial palpi with a distinct tuft at the apex beneath. Head and palpi silvery white. Second joint of the labial palpi brownish-gray, except at the tip. Antennae, thorax and anterior wings pale brownish-gray, in some lights silvery or golden. Thorax with a longitudinal median white streak produced gradually, widening along the dorsal margin of the anterior wings to the basal third, when it is intersected by a somewhat oblique costal white streak. Another oblique costal white streak about the basal fourth; a third one longer and narrower, situated beyond the middle of the costa, and a fourth and large one near the apex, and extending into the ciliae. Opposite the space between the second and third costal streaks is a large white triangular dorsal spot with its internal margin convex. Each of the costal and dorsal streaks is dark margined on both sides and around the apex. In the apex behind the costo-apical white spot is a curved golden-brown streak bordered behind by a brilliant white streak on the dorso-apical margin. Ciliae white, stained at the extreme tip with fuscous, with a rather wide, short, reddish-golden, hinder marginal line at the base of the apical ciliae, and three or four minute fuscous spots in the ciliae beyond it. Posterior wings bluish smoky. Legs golden-brown, spotted and streaked with white. Tarsi white, annulate with brown; posterior tibiae white. *Alar ex.* scarcely $\frac{1}{3}$ inch. Kentucky.

6. *Gracillaria salicifoliella*. N. sp.

Face and palpi white, with the apex of the labial palpi and a spot on the outer surface of the second joint, and one on the third, brown. Vertex white, suffused with brownish in front, and with a blackish patch at the base of the antennae, which are dark brown. Thorax and dorsal portion of the wings to the ciliae, white; costal portion blackish-brown; its line of junction with the white portion *twice indented or scalloped towards the fold*. Five costal white streaks, the first about the middle of the costa, curving backwards, long, and narrowed towards its apex, sometimes