

own accord. After two weeks' treatment there was very decided improvement, the bluish tint having faded considerably; some of the pustules had already disappeared. By the end of three months all the pustules were gone, and the erythema had disappeared likewise.

*Case 2.*—Mrs. R., thirty-six years of age, married, related a history of having had at various times, tonsillitis, neuralgia, acute Bright's disease after confinement. Five years previous to presenting for examination and treatment, she began to be troubled with dyspepsia, pain two or three hours after eating, gaseous eructations, and sour-brash. The pulse was normal, tongue clean and moist, appetite then good; heart normal. About the time the indigestion commenced five years ago, a small "boil" appeared at the right side of the nose near the commencement of the naso-labial fold. After that she never was free from large pimples in that region where five or six were always aggregated. It was so offensive to herself that she remained indoors a great deal rather than appear upon the street with such a disfiguring condition. There was one fairly large pustule between the eyebrows implanted upon a rosacea as was the patch at the side of the nose. They had been on the chin but had disappeared spontaneously. The ala nasi of the right side presented a deep flush, whilst the organ appeared broadened and slightly swollen at the tip. Otherwise there was no evidence of ill-health, and no history of alcoholism. Constipation had also been troublesome for a similar period of five years. This case cleared up rapidly upon treatment and was quite restored to the normal in four weeks' time.

*Case 3.*—Miss O. F., twenty years of age, single. She stated she had typhoid fever at four years; never had any indigestion; chicken-pox, the only disease of childhood. Had been troubled with constipation for two or three years; menstruation slightly irregular, as it sometimes lasted seven days, and the discharge very often scanty. She presented the appearance of good health. For four or five years she has had a most unsightly chin. The affection covered the whole skin surface between the vermilion border of the lower lip and the mental process, slanting downwards and outwards from the corners of the mouth. What skin could be seen was deeply stained, and there was not a space where the end of a match could be placed without touching a large yellow pustule or nodule. There was also slight erythema on the tip of the nose. The condition cleared up surprisingly fast under treatment, and by the end of a month there was only one pustule in the middle of the chin, but still a deep rosaceous background.