

of affairs exists. A very limited list of remedies of demonstrated value is presented for selection, and I believe I am not wide of the mark in saying that, in the hands of most practitioners, no remedy or combination of remedies hitherto in general use has been productive of anything but disappointment.

Some time ago my attention was drawn to Ergoapiol (Smith) as a combination of value in the treatment of a great variety of uterine disorders. Its exhibition in several cases in my hands yielded such happy results that I have used it repeatedly in a considerable variety of conditions, and with such uniformly good results that I am confirmed in the opinion that its introduction to the profession marks an era in modern therapeutics. In the treatment of irregular menstruation and attendant conditions I have found it superior to any other emmenagogue with which I am familiar, in the following important particulars:

1. It is prompt and certain in its action.
2. It is not nauseating and is not rejected by delicate stomachs.
3. It is absolutely innocuous.
4. It occasions no important after-effects.
5. It is convenient to dispense and administer.

The following clinical notes will afford a general idea of its action in a variety of cases:

CASE I.—Mrs. — came to me presenting the following symptoms incident to a delayed menstruation: Persistent headache of a neuralgic character; dull, aching pain in limbs and lumbar region; cramp-like pains in abdomen, and considerable nausea. The menstrual period was overdue seven days, but as yet there was no appearance of flow. Her periods had always been occasions of intense suffering, but had never before been delayed. I began the use of Ergoapiol (Smith), with some misgiving owing to the irritable condition of the stomach. One capsule every three hours was administered without any aggravation of the gastric distress. In twenty hours a normal menstruation was well under way; the flow was slightly increased over that observed on former occasions. The pains had subsided. Ergoapiol (Smith) was administered, one capsule three times a day, during the menstrual period, which terminated in five days. The patient was instructed to return for a quantity of the remedy several days before the next menstrual period. She did so, and, following directions, took one capsule three times a day for three days before expected menstruation. She subsequently reported that during the period—lasting five days—there had been practically no pain, and that the amount of flow was, as far as she could judge, normal.