

Children already wearing glasses should be treated with such glasses properly adjusted on the face.

Ascertain if the pupil habitually suffers from inflamed lids or eyes.

Place the testing card on the wall in a good light; do not allow the face of the card to be covered with glass.

The line marked xx (20) should be seen at 20 feet, therefore place the pupil twenty feet from the card.

Each eye should be examined separately.

Hold a card over one eye while the other is being examined. Do not press upon the covered eye, as the pressure might induce an incorrect examination.

Have the pupil begin at the top of the test card and read aloud as far as he can, first with one eye and then with the other.

If the pupil does not habitually suffer from inflamed lids or eyes, or can read a *majority* of the xx (20) test type with each eye, and does not, upon inquiry, complain of *habitually* tired painful eyes and headache after study, his eyes may be considered satisfactory. But, if he habitually suffers from inflamed lids or eyes or cannot read a *majority* of the xx (20) test type with both eyes, or habitually complains of tired or painful eyes or headache after study, a card of information should be sent to the parent or guardian.

FACTS TO BE ASCERTAINED.

1. Does the pupil habitually suffer from inflamed lids or eyes?
2. Does the pupil fail to read a majority of the letters in the number xx (20) line of the Snellen's Test Types, with either eye?
3. Do the eyes and head habitually grow weary and painful after study?
4. Is the pupil probably "cross-eyed"?
5. Does the pupil complain of earache in either ear?