- It seems to me that neither this nor the next present sufficient characters to warrant their being separated from *L. Canadensis* as anything more than varieties.
- 1311. L. HIRSUTA, Muhl.
 - L. Canadensis var. sanguinea, T. & G.
 - Railway banks and rocky woods, not uncorumon. Leaves all runcinate-pinnatifid, hirsute on the mid-ribs beneath. Flowers purplish-yellow. Aug.-2.
- 1314. L. LEUCOPHÆA, Gray.
 - Mulgedium leucophæum, D. C.
 - Low, rich wouls. A tall plant much like L. Canadensis, but with brownish pappus. Aug.-2.
- 1315. L. sativa, L. (Garden Lettuce.)
 - Occasionally found on waste heaps; but never permanently established.
- PRENANTHES, L. Rattlesnake-root.
 - 1316. P. ALBA, L. (White Lettuce.)

Nabalus albus, Hook.

- Rocky woods. An interesting plant with glaucous foliage and purplish-red stems, peduncles, and involucres. Flowers white and pappus deep brown. Aug.—2. (B.)
- 1318. P. ALTISSIMA, I.
 - Nabalus altissimus, Hcok.
 - Woods. A tall, slender plant with yellow flowers and dusky pappus. Aug.-2. (B.)
- SONCHUS, L. (Sow Thistle.)
 - 1321. S. oleraceus, L. (Common Sow-thistle.)
 - Cultivated ground and waste places. Annual. Aug.-1. Leaves clasping by a heart-shaped base, the auricles acute.
 - 1322. S. asper, Vill. (Prickly Sow-thistle.)
 - With the last, but commoner. Leaves more prickly, the auricles of the base rounded.
 - 1323. S. arvensis, L. (Perrennial Sow-thistle.)
 - At the base of Parliament Hill is a large and rapidly increasing patch of this plant, which is now becoming a troublesome weed in Canada. It has also been noticed at Chelsea and at Billings Bridge at the edges of fields. Aug.—1.
- TRAGOPOGON, L. Goat's-beard.
 - 1:26. T. pratensis, L. (Yellow Goat's-beard.)
 - Along the railway near the St. Louis Dam. A few plants have been found at the above locality for the last 12 years. Flowers, large, yellow.
 - 1327. T. porrifolius, L. (Salsify.)
 - An escape from cultivation. In a hay-field at Billings Bridge and occasional on waste heaps. Flowers purple.