

*Tr. Chirata.*—Chiretta 2½ oz.; proof spirit 20 fl. oz. (Dose one to two drachms.) This is a tincture from the D. and remains unchanged.

*Tr. Cinchona.*—Yellow bark 4 oz.; proof spirit 20 fl. oz. (Dose one to three drachms.) The L. and E. were 4½ oz. and 60 grs. av. to the 20 fl. oz.

*Tr. Cinchona Composita.*—Pale bark 2 oz.; bitter-orange peel 1 oz.; serpentary ½ oz.; saffron 60 grs.; cochineal 30 grs.; proof spirit 20 fl. oz. (Dose one to three drachms.) The L. and E. put in 85 grs. more of the bark to the 20 oz.; otherwise this recipe differs but slightly from that of the three colleges.

*Tr. Cinnamomi.*—Cinnamon 2½ oz.; proof spirit 20 fl. oz. (Dose one to three drachms.) The L. and E. were scarcely 2 oz. to the 20 oz.

*Tr. Cinnamomi Co.*—This, although it occupied a place in the three pharmacopœias, has not been deemed worthy of a place in the new one.

*Tr. Cocci.*—Pulv. Cochineal 2½ oz.; proof spirit 20 fl. oz. (Dose from one to three drachms.) The D. was 2 oz. to 30. Used chiefly for colouring mixtures.

*Tr. Colchici Semina.*—Bruised colchicum seed 2½ oz.; proof spirit 20 fl. oz. (Dose from one to two drachms.) This was the D. formula, the L. and E. were 2½ oz. av. to 20 fl. oz.

*Tr. Colchici Comp.*—This has been omitted by the Council.

*Tr. Conii Fructus.*—Dried ripe hemlock fruit, bruised, 2½ oz.; proof spirit 20 fl. oz. (Dose from 15 to 30 m.) The fruit of the conium has been selected as being more certain in its strength than the dried leaves; the tincture of the latter has therefore been left out of the new work.

*Tr. Croci.*—Saffron 1 oz.; proof spirit 20 fl. oz. (Dose from one to three drachms.) This is the D. formula; the E. had 42½ grs. more in the 20 oz. Used as a colouring for mixtures.

*Tr. Cubeba.*—This has not been given a place, on account of the alcohol necessary for its preparation. We regret it, as the L. preparation was an excellent one, and must still continue to be employed.

*Tr. Cuparise.*—Was discarded as unnecessary.

*Tr. Digitalis.*—Foxglove leaves 2½ oz.; proof spirit 20 fl. oz. (Dose from 15 to 30 or 40 m, cautiously increased. In delirium tremens half an ounce.) This is the D. formula, the L. and E. being but 2 oz. 85 grs. av. (134 grs. less) to the 20 fl. oz.

*Tr. Ergote.*—Bruised ergot 5 oz.; proof spirit 20 fl. oz. (Dose from one to two drachms.) The D. was 4 oz. to 20.

*Tr. Ergota Eth.*—Of the L. P., was not adopted.

*Tr. Ferri Perchloridi.*—Iron wire 1 oz.; hydrochloric acid 5 fl. oz.; nitric acid 3 drachms; water 3½ oz.; alcohol 15 fl. oz. Dilute the muriatic acid with 2½ oz. of the water, add the wire and dissolve with a gentle heat in a glass or enamelled vessel. Next mix the nitric acid and the remaining ounce of water, then pour them into the solution of iron, and evaporate to 5 fl. ounces; when cold add the alcohol. (Dose 10 to 30 m. gradually increased to a drachm or more.)

The D. was prepared in a similar manner but contained 3½ oz. av. of wire in 20 fl. oz. and was consequently 3½ times as strong. The other colleges directed the oxide of iron to be dissolved in the hydrochloric acid; the product was of the same strength as that of the new pharmacopœia, (viz., of the sp. gr. 0.992).

*Tr. Galla.*—Bruised galls. 2½ oz.; proof spirit 20 fl. oz. (Dose from half to two drachms.) This is the D. receipt, the L. and E. were 2½ oz. av. to 20 fl. oz.

*Tr. Gentiane Co.*—Gentian 1½ oz.; bitter-orange peel ½ oz.; cardamoms ½ oz.; proof spirit 20 fl. oz. (Dose from half to two drachms.) This is the D. receipt, the L. and E. contained 5½ grs. less of gentian in the 20 oz.

*Tr. Guaiaci.*—This tincture has been omitted.

*Tr. Guaiaci Ammoniata.*—Guaiac resin 4 oz.; aromatic spits. ammonia 20 fl. oz. (Dose one to two drachms in linseed tea or gruel.) The L. and E. were 3½ oz. and 40 grs. av. to 20 fl. oz.

*Tr. Hellebori.*—The tincture of black hellebore has not been thought worthy of a place in the new pharmacopœia.

*Tr. Hyoscyami.*—Dried henbane leaves 2½ oz.; proof spirit 20 fl. oz. (Dose one fluid drachm.) This is the D. recipe, the L. and E. being nearly 2½ oz. to the 20 fl. oz.

*Tr. Iodi.*—Iodine ½ oz.; iodide of potassium ½ oz.; alcohol 20 fl. oz. (Dose 15 to 30 m. to be gradually increased.) This tincture is scarcely strong enough for external application for which another preparation is given under the name of Linimentum Iodi. Of this former tincture the L. was iodine ½ oz. and 21 grs.; and iodide of potassium 1 oz. and 4½ grs. to 20 fl. oz. The E. was iodine 1½ oz. and 53 grs. av. to 20 fl. oz. alcohol. The D. was iodine ½ oz.; iodide of potassium 1 oz.; alcohol 20 fl. oz.

*Tr. Jalape.*—Jalap 2½ oz.; proof spirit 20 fl. oz. (Dose from one to three drachms.) This is the D. recipe. The L. was 2½ oz. to 20 fl. oz. The E. was 3½ oz. and 39 grs. to 20 fl. oz.

*Tr. Kino.*—Kino 2 oz.; alcohol 20 fl. oz. (Dose from one to two drachms.) The L. and E. had 3½ grains less in the 20 fl. oz.

*Tr. Kramerie.*—Rhatany 2½ oz.; proof spirit 20 fl. oz. (Dose from one to two drachms.) The D. was 4 oz. to 20 fl. oz.

*Tr. Lavandule Co.*—Oil of lavender 45 m.; oil of rosemary 5 m.; cinnamon bruised 75 grs.; nutmegs 75 grs.; red saunders 150 grs.; alcohol 20 fl. oz. (Dose 30 drops to a drachm, in water or in sugar.) This is a copy of the L. formula, the D. was made from the distilled spirits but did not vary greatly in strength. The D. was about twice as strong as the above.

*Tr. Limonis.*—Fresh lemon peel 2½ oz.; proof spirit 20 fl. oz. (Dose one or two drachms.) The L. was scarcely 2 oz. av., whilst the D. put 5 oz. to the 20 fl. oz.

*Tr. Lobelie.*—Lobelia herb 2½ oz.; proof spirit 20 fl. oz. (Dose one or two drachms; emetic dose ½ oz.) This was the D. recipe, the L. and E. were very nearly 2½ oz. to the 20 oz. proof spirit.

*Tr. Lobelie Ethera.*—Lobelia herb dried 2½ oz.; sulphuric ether 6½ fl. oz.; alcohol 13½ fl. oz. (Dose one or two drachms.) The L. and E. contained nearly ½ oz. more of the herb to the 20 oz.

*Tr. Lupuli.*—Hops 2½ oz.; proof spirit 20 fl. oz. (Dose one to two drachms in syrup.) The L. was a little over 3½ oz., hops to 20 fl. oz. The E. and D. employed the yellow powder (lupulin) sifted out of the hops, of which the E. directed 2½ oz. and the D. 2½ oz. to the Imperial pint of proof spirit. The hops themselves make an uncertain tincture. It is to be regretted that the compilers were not aware of this fact.

*Tr. Matico.*—This has not found a place in the new work.