

the use of gradually increasing doses of Bovinine, in milk by the stomach ; combined with complete rest in bed and forced feeding. I have had two cases of ulcer of the stomach, where the hemorrhage was severe, and collapse imminent, yield quickly to high rectal injections, of ounce doses of Bovinine and salt water, every three hours. I placed both of these patients upon glauher salts to procure an alkalinity of the blood, and to overcome constipation, and when the stomach was able to bear nourishment, I gave teaspoonful doses of pure Bovinine, to begin, and gradually increased the dose to four times daily. This treatment, combined with a proper diet, and rest in bed, brought about a complete cure in both cases.

In conclusion, gentlemen, I might say, that I am very much averse to recommending a proprietary preparation for fear of being classed with those physicians who immediately rush into print, lauding to the skies each new "cureall" that is put upon the market. On the other hand, I do not believe that a physician violates the code of ethics when he comes out squarely for a reputable preparation that has stood him in good stead, as it would be unjust and cowardly for him to use it secretly. Some of you may ask, If blood is what is needed in these cases, why not use it directly? Why depend on a proprietary article? In answer to these questions I would say that I have not the necessary knowledge, time or chemical appliances necessary to elaborate the arterial blood of the ox ; no more than I have the knowledge. etc., to make my morphine from the poppy. Therefore, in the cases where Haematherapy is indicated, I have always used, and shall continue to use, Bovinine.